

Important information. Please read carefully

Sedilix[®]-DM

Linctus

For the relief of resistant, dry and unproductive cough

Composition

Each 5ml linctus contains:

Dextromethorphan Hydrobromide 15mg
Pseudoephedrine Hydrochloride 30mg
Promethazine Hydrochloride.....3.125mg
Parabens 0.125% w/v as preservative

Pharmacodynamics

Dextromethorphan is a non-narcotic cough suppressant which acts centrally to raise the threshold for coughing. Its effectiveness in suppressing cough has been found to be about equal to codeine. Pseudoephedrine hydrochloride is used as a nasal and bronchial decongestant. It is said to have less pressor activity and CNS effects than ephedrine of which it is a stereoisomer. Promethazine hydrochloride is a H₁-receptor blocking agent. In therapeutic dosages, promethazine produces no significant effects on the cardiovascular system.

Pharmacokinetics

Dextromethorphan is rapidly absorbed from the gastrointestinal tract. It is metabolised in the liver and excreted in the urine as unchanged dextromethorphan and demethylated metabolites including dextrorphan, which has some cough suppressant activity.

Pseudoephedrine is readily absorbed from the gastrointestinal tract. It is excreted largely unchanged in the urine with small amounts of its hepatic metabolite. It has a half-life of about 5 to 8 hours; elimination is enhanced and half-life accordingly shorter in acid urine. Small amounts are distributed into breast milk.

Promethazine is well absorbed after oral or intramuscular doses. Peak plasma concentrations have been seen 2 to 3 hours after a dose by these routes, although there is low systemic bioavailability after oral doses, due to high first-pass metabolism in the liver. Promethazine crosses the blood-brain barrier and the placenta, and is distributed into breast milk. Values ranging from 76 to 93% have been reported for plasma-protein binding. Promethazine undergoes extensive metabolism, predominantly to promethazine sulfoxide, and also to *N*-desmethylpromethazine. It is excreted slowly via the urine and bile, chiefly as metabolites. Elimination half-lives of 5 to 14 hours have been reported.

Indications

Sedilix-DM is indicated for the relief of dry irritating coughs such as those associated with common cold or upper respiratory tract infections.

Sedilix-DM helps to relieve nasal congestion, sneezing and rhinorrhoea associated with allergy and the common cold.

Dosage and Administration

- **Adults and children over 12 years:**
10ml every 6 hours;
- **Children 6 to 12 years:**
5ml every 6 hours;
- **Children 2 to 5 years:**
2.5ml every 6 hours;

Contraindications

Sedilix-DM should not be taken for persistent or chronic cough (eg with smoking, emphysema, asthma) or when coughing is accompanied by excessive secretions unless directed by a physician.

It should be avoided in patients with liver disease or asthmatic patients and is contraindicated in patients taking monoamine oxidase inhibitors and within 2 weeks from stopping treatment, known hypersensitivity to pseudoephedrine and during acute attacks of asthma.

Warnings and Precautions

This preparation may cause drowsiness. If affected, do not drive motor vehicle or operate machinery. Avoid alcoholic drink while on this medication.

Use with caution in patients with epilepsy, prostatic hypertrophy, glaucoma, hepatic disease, hypertension, heart disease, diabetes, hyperthyroidism, patients with stenosing peptic ulcer, pyloro-duodenal obstruction or bladder neck obstruction, unless under the medical advice and supervision.

If symptoms do not improve within one week or accompanied with high fever, consult a physician before continuing use.

As with any other drugs, use in pregnancy is best avoided.

This product contains promethazine hydrochloride. It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.

To be used with cautions, and as advised by the physician/pharmacist for children age 2 to 6 years.

This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Side Effects

Sedilix-DM may cause drowsiness, dizziness and constipation.

Other side effects that may occur include GIT discomfort. No apparent evidence of physical dependence of the morphine type.

Other less common side-effects may include transient hypertension, dry mouth, insomnia, restlessness, palpitations, allergic reactions such as rashes, tightness of chest, thickening of bronchial secretions, toxic psychosis and blood dyscrasia.

Drug Interactions

Possible reactions systemically may occur with betahistine, certain hypnotic and anxiolytic medication, metoclopramide and antiparkinson drugs.

Incompatibilities

This preparation is incompatible in solution with aminophylline, barbiturates, benzylpenicillin salts, carbenicillin and heparin.

Symptoms and treatment for overdose

In cases of overdose, hospital admission is strongly advised. Overdose may produce respiratory depression, paranoid psychosis, delusions, hallucinations, and convulsion.

Treatment should include emptying the stomach by aspiration or gastric lavage. Nervous stimulation and convulsions should be treated with a sedative such as diazepam intramuscularly. If marked excitation is present, a sedative such as diazepam or a short-acting barbiturate may be given.

Elimination of pseudoephedrine may be assisted by acidification of the urine with ammonium chloride. Other symptomatic and supportive treatments should be given.

Shelf-life

The expiry date is indicated on the packaging.

Presentation

Brown syrup in packs of 90ml and 120ml.

Storage

Store in a cool place below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 08-Mar-2019

Manufactured and Marketed by
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate
75250 Melaka, Malaysia.

PATIENT INFORMATION LEAFLET

Sedilix®-DM

Linctus

Please read this leaflet carefully before you start to take your medicine. This leaflet is a summary of important information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

What is in Sedilix®-DM Linctus?

The active ingredients of Sedilix®-DM Linctus are Dextromethorphan Hydrobromide, Pseudoephedrine Hydrochloride and Promethazine Hydrochloride.

Sedilix®-DM Linctus contains Dextromethorphan Hydrobromide 15mg, Pseudoephedrine Hydrochloride 30mg and Promethazine Hydrochloride 3.125mg in each 5ml of syrup. Also contains Parabens 0.125%w/v as preservative.

Dextromethorphan is a cough suppressant, Pseudoephedrine is a nasal decongestant and Promethazine is an antihistamine. Sedilix®-DM Linctus is a cough and cold mixture that used for the relief of resistant, dry and unproductive cough.

Sedilix®-DM Linctus is a brown clear syrup with cherry flavour. This syrup liquid is available in pack sizes of 90ml and 120ml.

What is Sedilix®-DM Linctus used for?

Sedilix®-DM Linctus is indicated for the relief of dry irritating coughs such as those associated with common cold or upper respiratory tract infections.

Sedilix®-DM Linctus helps to relieve nasal congestion, sneezing and rhinorrhoea associated with allergy and the common cold.

Sedilix®-DM Linctus is for short term use only. If symptoms persist longer than 7 days or accompanied by fever, consult a physician.

How much and how often should you use Sedilix®-DM Linctus?

Adults and children above 12 years: 10ml every 6 hours;

Children 6 to 12 years: 5ml every 6 hours

Children 2 to 5 years: 2.5ml every 6 hours

Extreme care should be taken in giving an accurate dose of medicine because overdoses have been reported to cause serious adverse events. It is highly recommended that a measuring cup, spoon or syringe (which can be obtained from the pharmacy) be used to administer dose.

When should you not take this medicine?

Sedilix®-DM Linctus should not be taken for persistent or chronic cough (eg with smoking, emphysema, asthma) or when coughing is accompanied by excessive secretions unless directed by a doctor.

Not for use in patient suffering from severe high blood pressure and severe coronary artery disease (heart disease caused by poor blood flow or narrowing of the blood vessels of the heart).

You should not use this medicine if you have high blood pressure, heart condition, angle closure glaucoma, prostate hypertrophy or hyperthyroidism (a thyroid disease). Please check with your doctor if you have any of the above conditions.

Sedilix®-DM Linctus should be avoided in patients with liver disease or asthmatic patients and is contraindicated in patients taking monoamine oxidase inhibitors or within 2 weeks from stopping such treatment known hypersensitivity to pseudoephedrine, promethazine, dextromethorphan and any ingredient of this product and during acute attacks of asthma.

As with all medicines, use in pregnancy is best avoided unless necessary, where close medical supervision is required. Do not take this medicine during pregnancy or breast feeding, before consulting your doctor or pharmacist.

Special Warnings

Sedilix®-DM Linctus is strictly not to be used for infants six months of age and younger.

It is also not recommended for children below 2 years old because younger children are more prone to serious side-effects of promethazine.

Sedilix®-DM Linctus is also to be used with caution in children 2 years and older.

To be used with cautions, and as advised by the physician/pharmacist for children age 2 to 6 years.

As the preparations may cause drowsiness, adult patients should be advised against driving or operating machinery and to avoid consuming alcoholic drinks while on this medication.

This medicine may cause your skin to be more sensitive to the sun especially if you have eczema or a tendency to rheumatism. You should protect your skin from exposure to strong sunlight.

If you are taking other medicines, going for a laboratory test or suffering from other medical conditions, consult your doctor or pharmacist before taking this medicine.

If symptoms do not improve within one week, or if symptoms are accompanied with high fever, consult your doctor before continuing use.

This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Does Sedilix®-DM Linctus have side effects?

A medicine may cause side effects in certain individuals. If you experience side effects and are concerned, you should seek advice from your doctor or pharmacist.

Sedilix® may cause drowsiness, dizziness and constipation.

Other side effects that may occur include stomach upset diarrhoea, loss of appetite, nausea or vomiting, blurred vision and tinnitus (ringing or buzzing in ears).

Other less common side-effects may include fever, breathing difficulty, jaundice (yellow tinge to skin or eyes), tremors or convulsions, transient hypertension, dry mouth, insomnia, restlessness, irregular heart beat, allergic reactions such as rashes, tightness of chest, thickening of bronchial secretions, toxic psychosis and blood dyscrasia.

What other medicine or food should be avoided whilst taking Sedilix®-DM Linctus?

Avoid using alcohol, sleeping pills, medicines for reducing anxiety and tension or any other medicines that suppress breathing.

What should you do if you miss a dose?

If you are taking this medicine regularly and miss a dose, take it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take a double dose to make up for the dose that you have missed.

How should you keep Sedilix®-DM Linctus?

Keep Sedilix®-DM Linctus below 30°C, in a safe place and out of reach of children.

What are the symptoms of overdose and what to do when you have taken more than the recommended dosage?

Symptoms of overdose include over-excitement, shaky and unsteady movements, convulsions and/or tremors, hallucinations, difficulty in breathing, drowsiness, unconscious, fixed dilated pupils with a flushed face and fever.

If you experience any of the symptoms, or if you have taken more than the recommended dose, seek immediate medical attention.

When should you consult your doctor?

You need to consult a doctor when:

- you experience common side effects that bother you and serious side effects that need urgent medical attention.
- you have taken this medicine for 7-10 days in a row and your condition is not improving.
- you have taken an overdose of the medicine.