



GUADEx LINCTUS

DESCRIPTION : Brown syrupy liquid, mixture of menthol, dill oil, peppermint oil & eucalyptus oil flavour and odour. Each 5ml contains dextromethorphan hydrobromide 15mg and guaiphenesin 100mg with 0.2% w/v sodium benzoate as preservative.

PHARMACODYNAMICS : Dextromethorphan hydrobromide is a d isomer of the codeine analog of levorphanol. The drug acts centrally to elevate the threshold for coughing. Guaiphenesin is reported to reduce the viscosity of the tenacious sputum and increase secretions of the respiratory tract.

PHARMACOKINETICS : Dextromethorphan hydrobromide acts centrally to elicit a cough suppressant response, and has marked antitussive properties of codeine except the analgesic properties and addictive properties of codeine. Dextromethorphan hydrobromide is well absorbed from the gastrointestinal tract. It is metabolised in the liver and excreted as unchanged dextromethorphan hydrobromide and demethylated morphine compound. Guaiphenesin is readily absorbed from the G.I tract and rapidly metabolised and excreted in the urine.

INDICATIONS : For dry non-productive cough only.

CONTRAINDICATIONS : Known hypersensitivity to any component in the formulation.

DRUG INTERACTIONS : Serious toxicity has been reported in patients receiving dextromethorphan hydrobromide and monoamine oxidase inhibitors (MAOIs), including 14 days after stopping the MAOIs drug. The use of the dextromethorphan hydrobromide may result in additive CNS depressant effects when coadministered with alcohol, antihistamines, psychotropics or other drugs which produce CNS depression.

ADVERSE EFFECTS : Dextromethorphan hydrobromide occasionally causes drowsiness, dizziness, excitation, mental confusion and gastrointestinal disturbances. Very high dose may produce respiratory depression. Guaiphenesin may produce gastro-intestinal discomfort and drowsiness. Very large dose may cause nausea and vomiting.

PRECAUTIONS AND WARNINGS : May cause drowsiness in some patients and those affected should be advised not to drive or operate machinery. It is dangerous to exceed the stated dose.

- i) Not to be used in children less than 2 years of age.
- ii) To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

PREGNANCY & LACTATION : Unknown.

RECOMMENDED DOSAGE : Adult : 5ml-10ml; Children 7 to 12 years : 5ml; 2 to 6 years : 2.5ml; Doses to be given undiluted 3 times a day.

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTE(S) : Dextromethorphan hydrobromide may elicit symptoms similar to codeine phosphate overdose which may result in coma, pulmonary oedema, respiratory depression and CNS excitation including convulsion.

Treatment:

1. Maintain respiratory functions, artificial ventilation and oxygen therapy.
2. Induce vomiting unless patient is comatose.
3. Naloxone may be administered as an antidote.

PACKING : 90ml PET amber bottle.

STORAGE : Keep container well closed. Protect from strong light. Store below 30°C. Keep out of reach of children. Recommended shelf-life : 3 years.

MANUFACTURED BY :

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohd. 6, Bandar Sultan Suleiman,
42000 Port Klang, MALAYSIA.

Revision date : June 2013

20032C