

LEAFLET



SW BROMHEXINE ELIXIR 4MG/5ML/ TABLET 8MG

	Bromhexine Elixir	Bromhexine Tablet
Bromhexine HCl	4 mg / 5ml	8 mg / Tablet
Preservative: Methyl Paraben	10 mg / 5ml	-

Pharmacology:
Bromhexine is a synthetic derivative of the herbal active ingredient vasicine. Preclinically, it has been shown to increase the proportion of serous bronchial secretion. Bromhexine enhances mucous transport by reducing mucous viscosity and by activating the ciliated epithelium (mucociliary clearance). In clinical studies, bromhexine showed a secretolytic and secretomotor effect in the bronchial tract area, which facilitates expectoration and eases cough.
Bromhexine hydrochloride is rapidly absorbed from the gastro-intestinal tract and about 85 to 90% of a dose is excreted in the urine mainly as metabolites. Ambroxol is a metabolite of bromhexine. Bromhexine is highly bound to plasma proteins.
Administration of bromhexine HCl by mouth to healthy subjects produced peak plasma concentrations after about 1 hour. Only small amounts were excreted unchanged in the urine with a half-life of about 6.5 hours.

Indications:
It is indicated in bronchitis and other respiratory conditions as an aid to expectoration.

Contraindications:
It is contraindicated in patients who is hypersensitive to it and should be given cautiously to patients with gastric ulceration.

Side Effects/Adverse Reactions:
Gastro-intestinal side effects may occur occasionally and a transient rise in serum amino-transferase values has been reported.
Immune System Disorders
Frequency not known: Anaphylactic reactions including anaphylactic shock.
Skin and Subcutaneous Skin Disorders
Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Precautions/Warnings:
Bromhexine should be given cautiously to patients with gastric ulceration. An increase in volume of liquefied sputum should be expected.
Very rare cases of chronically associated severe skin impairments such as Stevens Johnson syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Usage in Pregnancy and Lactation:
The safe use of Bromhexine HCl has not been established with respect to the possible adverse effects upon foetal development. Therefore, it should not be used in women of child bearing potential, particularly during early pregnancy, or in lactating women unless advised by the physician.

Drug Interactions:
Administration of Bromhexine together with Erythromycin or Cephalixin may enhanced this antibiotic concentration in bronchial secretion / lung tissue.

Dosage:
Oral administration.
Bromhexine Elixir 4mg/5ml:
Children less than 2 years : 1mg three times daily.
Children 2 to <6 years : 2mg three times daily.
Children 6 to <12 years : 4mg three times daily.
Adult : 8 to 16mg three times daily.

Bromhexine Tablet 8mg:
Children 6 to <12 years : 4mg three times daily.
Adult : 8 to 16mg three times daily.

Symptoms and Treatment of Overdose:
No symptoms of overdosage have been reported in man to date. If they occur, symptomatic treatment should be provided.

Pack Size: Elixir : A bottle of 60ml, 100ml and 120ml.
Tablet : A bottle of 50 tablets.
Blister pack : A box of 10 x 10, 50 x 10 and 100 x 10 tablets per strip.
Pack Size (export only): Elixir : A bottle of 1 litre.
Tablet : A bottle of 1000 tablets.

Storage Conditions: Store at or below 30°C. Protect from light.
Shelf-life: Elixir - 2 years Tablet - 3 years

Description: Bromhexine Elixir : A clear, colourless syrup with cherry and vanilla flavour.
Bromhexine Tablet : Pentagon shaped, white, flat, plain tablet with single score on one side only.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN/LOCAL GENERAL PRACTITIONER/PHARMACIST.

Manufacturer & Product Registration Holder:
Malaysia:
SUNWARD PHARMACEUTICAL SDN. BHD.
No. 9 ,11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

Singapore:
SUNWARD PHARMACEUTICAL PTE. LTD.
11, Wan Lee Road, Singapore 627943

CL305A-R13
Revision Date: 16/12/2024