

PACKAGE INSERT

ONADRINE TABLET

Each tablet contains:
Orphenadrine Citrate 50mg

Product Description: Round, white, standard convex, plain tablet with score on one side.

Pharmacodynamics:

Orphenadrine Citrate is a centrally acting (brain stem) compound, which in animals selectively blocks facilitatory functions of the reticular formation. Orphenadrine does not produce myoneural block, nor does it affect crossed extensor reflexes. Orphenadrine prevents nicotine-induced convulsions but not those produced by strychnine. The mode of therapeutic action of Orphenadrine has not been clearly identified, but may be related to its analgesic properties. Orphenadrine citrate also possesses anticholinergic actions.

Pharmacokinetics:

Orphenadrine is readily absorbed from the gastrointestinal tract and is almost completely metabolised to at least 8 metabolites. It is mainly excreted in the urine as metabolites and unchanged drug. A half life of 14 hours has been reported.

Indication:

Orphenadrine Citrate is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute painful musculo-skeletal conditions.

Recommended Dosage:

Two tablets daily, one in the morning and one in the evening.

Route of Administration: Oral

Contraindications:

Orphenadrine Citrate is contraindicated in patients with glaucoma, pyloric or duodenal obstruction, stenosing peptic ulcers, prostatic hypertrophy or obstruction of the bladder neck, cardio-spasm (megaesophagus) and myasthenia gravis.

Warnings and Precautions:

Orphenadrine Citrate should be used with caution in patients with tachycardia, cardiac decompensation, coronary insufficiency or cardiac arrhythmias. Some patients may experience transient episodes of dizziness or syncope, which may impair their ability to engage in potentially hazardous activities such as operating machinery or driving a motor vehicle. As with any drug, patients should be advised against the ingestion of alcohol during treatment. Orphenadrine has been shown to induce enzymatic systems involving the metabolism of aminopyrine, steroidal contraceptives, griseofulvin, hexobarbital and phenylbutazone. Orphenadrine will potentiate other anticholinergic agents.

Interactions with Other Medicaments:

Orphenadrine has been shown to induce enzymatic systems involving the metabolism of aminopyrine, steroidal contraceptives, griseofulvin, hexobarbital and phenylbutazone. Orphenadrine will potentiate other anticholinergic agents.

Pregnancy and Lactation:

As there are no adequate and well-controlled studies in pregnant women, this drug should be used only if clearly needed. The use of this drug in late pregnancy is not recommended because of the effects on the foetal cardiovascular system. Since safety of the use of this preparation in pregnancy, during lactation, or in the childbearing age has not been established, use of the drug in such patients requires that the potential benefits of the drug be weighed against its possible hazard to the mother and child.

Side Effects:

Adverse effects of orphenadrine are mainly due to the mild anticholinergic action of orphenadrine, and are usually associated with higher dosage. Dryness of the mouth is usually the first adverse effect to appear when the daily dose is increased. Possible adverse effects includes; tachycardia, palpitation, urinary hesitancy or retention, blurred vision, dilatation of pupils, increased ocular tension, weakness, nausea, vomiting, headache, dizziness, constipation, drowsiness, hypersensitive reactions, pruritus, hallucinations, agitation, tremor, gastric irritation, and rarely urticaria and other dermatoses.

Symptoms and Treatment of Overdose:

The symptoms of Onadrine includes :-
Convulsion, cardiac dysrhythmias, dry mouth, palpitation, urinary retention, blurred vision, dilatation of pupils, increase ocular tension, weakness, nausea, vomiting, headache, dizziness, drowsiness, hallucination, agitation and gastric irritation. If the overdosage occur, the stomach should be emptied by aspiration and lavage followed by the administration of active charcoal. Convulsion may be controlled by tubocurarine and intermittent positive – pressure ventilation. Excitement may be controlled by diazepam or a short-acting barbiturate. Supportive therapy may require oxygen and assisted respiration, ice-bag or alcohol sponges for hyperpyrexia, especially in children, bladder catheterisation and the administration of fluids.

Effects on Ability to Drive and Use Machine: Not applicable.

Storage Condition: Store at or below 30°C. Protect from light.

Pack Size: Blister Pack: A box of 10x10, 50x10 and 100x10 tablets per strip.

Pack Size (Export to Singapore): A bottle of 1000 tablets.

Shelf-life: 3 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder
SUNWARD PHARMACEUTICAL SDN. BHD.
No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia

TL654A-R6
Revision Date: 28/10/2025