

VENTAMOL EXPECTORANT

VIVEN02-var

DESCRIPTION

Strawberry flavoured, colourless, transparent liquid.

COMPOSITION

Each 5 mL contains:

Salbutamol (as sulphate)	1.2 mg
Guaiphenesin	50.0 mg

PHARMACODYNAMICS

Salbutamol is a direct-acting sympathomimetic amine. It acts relatively selectively on beta-2 adrenergic receptors in the lungs to relax bronchial smooth muscle, thereby relieving bronchospasm and reducing airway resistance.

Guaiphenesin has an expectorant action which increases the output of respiratory tract fluid by reducing adhesiveness and surface tension. The increased flow of less viscid secretions promotes ciliary action and facilitates the removal of mucus. This changes a dry, unproductive cough to a cough that is productive and less frequent.

PHARMACOKINETICS

Salbutamol and guaiphenesin are readily absorbed from the gastrointestinal tract. Major route of excretion is via renal system.

INDICATIONS

- To reduce the viscosity of tenacious sputum.
- To reduce bronchospasm and airway resistance in asthmatic patients.

CONTRAINDICATIONS

- Should not be used in hypersensitive patients.
- Ventamol Expectorant and beta-blocking drugs such as propranolol should not be prescribed together.

WARNINGS AND PRECAUTIONS

- Use of this medication should be carefully considered when the following medical problems exist: cardiovascular disorders, including coronary insufficiency and hypertension; diabetes mellitus; hyperthyroidism; thyrotoxicosis.
- Tocolysis: Serious adverse reactions including death have been reported after administration of terbutaline / salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established.

Adequate and well-controlled studies in humans have not been done.

It is not known whether salbutamol and guaiphenesin is distributed into breast milk of humans.

DRUG INTERACTIONS

- Concurrent use with corticosteroids may exacerbate the adverse effect of salbutamol.
- Propranolol and other beta-blockers antagonise the effect of salbutamol.
- Concomitant administration of aminophylline or other xanthines may enhance the effect of salbutamol.

MAIN SIDE/ADVERSE EFFECTS

Nausea, nervousness, trembling, unusual fast or pounding heartbeat, drowsiness, diarrhoea, stomach pain.

OVERDOSE AND TREATMENT

Symptoms: Tremor, bursting feeling, palpitations, agitation, tachycardia, peripheral vasodilatation, chest pain, hypokalaemia, hyperglycaemia, lactic acidosis, acute psychosis.

Treatment: By emesis or gastric lavage, if appropriate; and by administration of atenolol 2.5 - 10 mg or propranolol 1 mg/min.

DOSAGE AND ADMINISTRATION

Adults

Oral, 10 to 20 mL two to three times a day.

Children

Up to 6 years: Oral, 5 to 10 mL two times a day.
7 to 12 years: Oral, 10 mL two times a day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 25°C. Protect from light.

Presentation/Packing:

Expectorant x 100 mL and 3.6 Litres.

(Not all presentation available locally)

Product Registration Holder / Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: July 2018

hovid