

LOBESOL CREAM 0.05% W/W

Composition:	0.05% w/w
Clobetasol Propionate	
Ethylene Diaminetetraacetate	
Chlorocresol	
Cetostearyl Alcohol	
Deionised Water	
White Soft Paraffin	
Propylene Glycol	
Cetomacrogol 1000	

Product Description:
A white and smooth cream.

Pharmacodynamics:
Clobetasol propionate is a highly active corticosteroid with topical anti-inflammatory activity. The major effect of clobetasol propionate on skin is a non-specific anti-inflammatory responses, as a result of vasoconstriction and decrease in collagen synthesis.

Pharmacokinetics:
Percutaneous penetration of clobetasol propionate varies among individuals and can be increased by the use of occlusive dressings, or when the skin is inflamed or diseased.
Mean peak plasma clobetasol propionate concentrations of 0.63 nanograms/ml occurred in one study eight hours after the second application (13 h after an initial application) of 30g clobetasol propionate 0.05% ointment to normal individuals with healthy skin. Following the application of a second dose of 30g clobetasol propionate cream 0.05% mean peak plasma concentration were slightly higher than the ointment and occurred 10 h after application. In a separate study, mean peak plasma concentration of approximately 2.3 nanograms/ml and 4.6 nanograms/ml occurred respectively in patients with psoriasis and eczema three hours after a single application of 25g clobetasol propionate ointment. Following percutaneous absorption of clobetasol propionate the drug probably follows the metabolic pathway of systemically administered corticosteroids, i.e. metabolised primarily by the liver and then excreted by the kidneys. However, systemic metabolism of clobetasol has never been fully characterised or quantified.

Indication:
It is used for the short-term relief of the inflammatory and pruritic manifestations of moderate to severe corticosteroid-responsive dermatoses.

Recommended Dosage:
Apply the cream to the affected areas twice a day, preferably in the morning and evening. Generally, the dosage should not exceed 50gm of clobetasol propionate 0.05% cream per week and duration of a course of Clobetasol therapy should not exceed 14 days.

Route of Administration: Topical

Contraindications:
It is contraindicated in patient who are hypersensitive to Clobetasol propionate or other corticosteroids.

Warnings and Precautions:
Clobetasol propionate is one of the most potent topical corticosteroid preparations. It can suppress the hypothalamic-pituitary-adrenal axis following topical application, and HPA axis suppression has occurred following topical dosages as low as 2gm of the 0.05% cream daily. Because of the drug's potency and potential for causing adverse systemic effect during topical therapy, the usual dosage should not be exceeded and occlusive dressings should not be applied to areas of Clobetasol propionate application.

Visual Disturbance
Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to

an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Interaction with Other Medicaments:
Co-administered drugs that can inhibit CYP3A4 (e.g. ritonavir and itraconazole) have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the CYP3A4 inhibitor.

Pregnancy and Lactation:
Pregnant women and nursing mother should seek the advice of a physician before using this product. Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development. The relevance of this finding to human beings has not been established; however, topical steroids should not be used extensively in pregnancy, i.e. large amounts or for prolonged periods.

Side Effects:
The side effects are itching, burning, stinging, dryness, folliculitis, hypertrichosis, acne form eruption, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, milaria. Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal axis suppression, manifestation of Cushing's syndrome, hyperglycemia, and glucosuria in some patients. Vision, blurred.

Symptoms and Treatment of Overdose:
Topically applied corticosteroids can be absorbed in sufficient amount to produce systemic effect. Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressings should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid. Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favourable response does not occur promptly, the corticosteroids should be discontinued until the infection has been adequately controlled.

Effects on Ability to Drive and Use Machine: Not applicable.

Storage Conditions: Store at or below 30°C.

Marketed in Malaysia:	Marketed in Singapore:
Pack Size: A tube of 15g and 30g.	Pack Size: A jar of 400g.
A box of 50 x 15g/tube.	A tube of 15g and 30g.
A jar of 100g.	
Pack Size (export only): A jar of 400g.	

Shelf-Life: 3 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder: Sunward Pharmaceutical Sdn. Bhd. No. 9, 11&17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia.	Imported by : Sunward Pharmaceutical Pte. Ltd. 11, Wan Lee Road, Singapore 627943.
---	--

EL211A-R9
Revision Date: 27/09/2022