

Rowatanal® Cream

Name and Strength of Active Substances

The active substances are menthol, bismuth subgallate, calcium carbonate and zinc oxide. Each 100g of cream contains 1g menthol, 2g bismuth subgallate, 5g calcium carbonate and 8g zinc oxide.

Other Ingredients

The other ingredients are: hard paraffin, glycerol, wool fat, glycerol mono-dioleate, polyoxyethylene-12-cetyl stearyl alcohol, purified water and as preservatives: butyl parahydroxybenzoate, ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216) and methyl parahydroxybenzoate (E218).

Product Description

Rowatanal® Cream is smooth in texture and light yellow in colour.

Pharmacodynamics/Pharmacokinetics

Rowatanal® Cream has antiseptic and astringent (cleansing) properties. It relieves the intense itching and burning sensation associated with haemorrhoids.

Indication

For the treatment of haemorrhoids and proctitis (anal irritation).

Recommended Dosage

Apply 2-3 times daily and after defecation. For internal haemorrhoids, use the cannula nozzle provided.

Route of Administration

For rectal use only.

Contraindications

Do not use Rowatanal® if you:

- are allergic to menthol, bismuth subgallate, calcium carbonate, zinc oxide or any of the other ingredients of this medicine.

Warnings and Precautions

Talk to your doctor or pharmacist before using Rowatanal® if your piles are bleeding or have been present for some time.

If the condition persists or worsens you should also consult your doctor.

Rowatanal® contains glycerol, wool fat and parahydroxybenzoates. This medicine contains glycerol, which may have a mild laxative effect. It contains wool fat, which may cause local skin reactions (e.g. contact dermatitis). It also contains parahydroxybenzoates which may cause allergic reactions (possibly delayed).

Effect on ability to Drive and Use Machines

Rowatanal[®] does not normally produce any effects on ability to drive or use machines.

Interaction with Other Medicaments

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Statement on Usage during Pregnancy and Lactation

This product should not be used during pregnancy and lactation unless considered essential by your physician.

Ask your doctor or pharmacist for advice before taking any medicine.

Adverse Effects/Undesirable Effects

Like all medicines, this medicine can cause side effects, although not everyone gets them.

Side effects have been reported rarely with use of this and similar products. Side effects include a brief cooling sensation on application of the cream. Sensitivity reactions have been reported very rarely.

Rowatanal[®] contains glycerol, wool fat and parahydroxybenzoates. (See Warnings and Precautions for further information).

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Storage Conditions

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on both the outer carton and on the tube.

The expiry date refers to the last day of that month.

Do not store above 30°C.

Dosage Forms and Packaging Available

Rowatanal[®] is available in a 26g tube.

A white cannula nozzle is also included.

Name and Address of Manufacturer

ROWA[®] Pharmaceuticals Ltd., Bantry, Co. Cork, Ireland.

Product Registration Holder

DCH Auriga (Malaysia) Sdn. Bhd., Lot 6, Persiaran Perusahaan, Seksyen 23, 40300 Shah Alam, Selangor.

Date of revision of PI

This leaflet was last revised in November 2018.

Artwork Checklist	Job Description		Version Number	Size	Inks			ACC#
	Rowatan Carbon Tube PIL RCF 10546				Block			
	Market		Proof Date	Cutter Ref.		Colour		
	Malaysia		28/11/2018	N/A		Front Back Stroke		