



○ SWKD

○ WKMS

○ WKMM

○ HU

○ HU (KK)

File Name : LemonSweetPurgative\_Leaflet\_08122020

Received : 15.01.2021

Amended : -

Output : -

Requestor : Shuying

PIC :

Designer : Nickles

Version : 00

Opened : 150w x 210 MM

Closed : -

Finished :

Colour : 1C

K

# Three Legs Lemon Sweet Purgative Ubat Sembelit Perisa Lemon Cap Kaki Tiga

## 三脚标柠檬润肠露



**Composition:** Each 5ml contains 1.77g Magnesium Sulphate, 0.055ml Glycerin BP and 5mg Sodium Benzoate

**Product Description:** Clear solution with a citrus lemon odour and slight bitter taste.

**Indication:** For the relief of constipation and difficulty in bowel movement due to hardened faeces.

**Dosage:** Adults: 8 teaspoonful (40ml)  
Children over 6 years: 2 - 5 teaspoonful (10-25ml)

To be taken once a day. To be mixed with a full glass (240ml) of water before drinking. To be taken an hour before food or on an empty stomach.

**Route of Administration:** Oral

**Contraindications:**

- Internal use is contraindicated in all cases of acute gastro- intestinal conditions (except constipation), renal impairment and in children with intestinal parasitic diseases.
- Hypersensitivity to magnesium sulphate.

**Warnings and Precautions:**

- Should not be given to children less than 6 years unless prescribed by a doctor.
- Magnesium Sulphate should be given with care to patients with impaired kidney function.

**Effects on Ability to Drive and Use Machines:** Not known

**Interactions with Other Medicaments:** Oral magnesium salts have the properties of antacids. Therefore, it is recommended that this product is not taken within two to four hours of any other medicinal products to minimize interaction.

**Pregnancy and Lactation:** Do not use in pregnancy or while breastfeeding.

**Adverse Effects/ Undesirable Effects:** Hypermagnesaemia may occur after prolonged usage of magnesium sulphate as a purgative. May cause colic. Ingestion of magnesium salts may cause gastrointestinal irritation and watery diarrhoea. Rarely paralytic ileus has been reported.

**Symptoms and Treatment of Overdose:** Symptoms of hypermagnesaemia may include extreme thirst, a feeling of heat, hypotension due to vasodilation, drowsiness, nausea, vomiting, gastrointestinal irritation and watery diarrhoea, flushing, confusion, slurred speech, double vision and muscle weakness, loss of tendon reflexes due to neuromuscular blockade, CNS and respiratory depression, cardiac arrhythmias (including bradycardia), coma and cardiac arrest.

Treatment of mild hypermagnesaemia is usually limited to restricting magnesium intake. In severe hypermagnesaemia, ventilator and circulatory support may be required. Slow intravenous injection of calcium gluconate (10 to 20ml of 10% calcium gluconate) is recommended to reverse the effects on cardiovascular and respiratory systems. If renal function is normal, adequate fluids should be given to promote renal magnesium clearance. This may be increased by the use of furosemide. Haemodialysis using a magnesium- free dialysis solution effectively removes magnesium and this may be necessary in patients with renal impairment, or for whom other methods prove ineffective.

**Pharmacodynamics:** Magnesium Sulphate is a saline purgative. Its action as a purgative is because it is not readily absorbed from the intestine. When taken by mouth in dilute solution, it reduces the normal absorption of water from the intestine tract and retains water in the intestinal lumen by osmosis forces. The bulky fluid contents distend the bowel active reflex peristalsis is excited, followed by evacuation of the contents of the intestine.

**Pharmacokinetics:** When a dilute solution of magnesium sulphate is taken orally, the absorption of water from the intestine is reduced and the bulky fluid contents distend the bowel. Active peristalsis is excited and evacuation of the contents of the intestine results.

Magnesium sulphate cause bowel evacuation normally within 2- 4 hours.

**Storage Conditions:** Store below 30°C. Keep out of reach of children.

**Shelf Life:** 3 years

**Dosage Form:** Liquid

**Packing Size:** 40ml, 60ml, 100ml and 120ml

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

**Komposisi:** Setiap 5ml mengandungi 1.77g Magnesium Sulfat, 0.055ml Glyserin BP dan 5mg Natrium Benzoat

**Produk Deskripsi:** Cecair yang jelas dengan bau limau sitrus dan sedikit rasa pahit.

**Indikasi:** Untuk mengatasi masalah sembelit dan melawaskan pembuangan najis yang disebabkan oleh najis yang keras.

**Cara Penggunaan:** Dewasa: 8 sudu teh (40ml)  
Kanak- kanak lebih 6 tahun: 2 - 5 sudu teh (10-25ml)

Diminum sekali sehari. Perlu dicampur dengan segelas air (240ml) sebelum minum. Diminum sejam sebelum makan atau semasa perut kosong.

**Laluan Pemberian Ubat:** Oral

**Kontraindikasi:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Amaran dan Perhatian:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Kesan ke atas Keupayaan Memandu dan Penggunaan Mesin:** Tidak diketahui

**Interaksi dengan ubat lain:** Sila rujuk kepada penerangan Bahasa Inggeris

**Kehamilan dan Penyusuan:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Kesan Sampingan:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Simptom dan Rawatan untuk Kelebihan Dos:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Farmakodinamik:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Farmakokinetik:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Keadaan Penyimpanan:** Simpan di bawah 30°C. Jauhkan daripada capaian kanak-kanak.

**Tempoh Hayat:** 3 tahun

**Dos:** Cecair

**Saiz Bungkusan:** 40ml, 60ml, 100ml and 120ml

Maklumat terkandung dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

**成份：**每5毫升含有1.77克 硫酸镁，0.055毫升甘油和5毫克苯甲酸钠

**产品外观：**带有柑橘柠檬气味和微苦味道的清澈液体。

**主治：**用于解除便秘及因粪便硬化而引起的排便困难。

**服量：**成人：8茶匙（40毫升）  
6岁以上小孩：2–5茶匙（10–25毫升）

每日服一次。与一杯水（240毫升）混合后服用。在饭前一个小时或空腹服用。

**给药途径：**内服

**禁忌：**请参阅英文版本。

**警告及注意：**请参阅英文版本。

**对驾驶和使用机器的能力的影响：**未知

**药物相互作用：**请参阅英文版本。

**怀孕和哺乳：**请参阅英文版本。

**副作用：**请参阅英文版本。

**药物过量症状及治疗：**请参阅英文版本。

**药效学：**请参阅英文版本。

**药代动力学：**请参阅英文版本。

**贮存：**储藏低于30°C。远离儿童。

**有效期：**3年

**剂型：**液体

**包装：**40毫升、60毫升、100毫升及120毫升

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。

Manufactured by Product Registration Holder  
**Syarikat Wen Ken Drug Sdn.Bhd.**  
24, Jalan Lambak, Taman Johor,  
81200 Johor Bahru, Johor, Malaysia.

Singapore Distributor:  
**Wen Ken Marketing (S) Pte. Ltd.**  
2, Alexandra Road, #02-08 Delta House,  
Singapore 159919.

Date of Revision: 15.01.2021  
...