



TUSS-DEX SYRUP

Description: Brown coloured, sour plum flavoured syrup.

Contents:

Each 5ml contains :	Tripolidine HCl	1.25mg	Methylparaben	0.1% w/v
	Pseudoephedrine HCl	30mg	Propylparaben	0.01% w/v
	Dextromethorphan HBr	15mg		

Pharmacology:

Tripolidine HCl, a potent antihistamine, provides a drying effect on the nasal mucosa by means of its antimuscarinic actions. Pseudoephedrine, a sympathomimetic amine, acts on alpha-adrenergic receptors in the mucosa of the respiratory tract to produce vasoconstriction which reduces the swelling associated with inflammation of the mucous membranes lining the nasal passage. Dextromethorphan, a cough suppressant, acts centrally to elevate the threshold for coughing. It is well absorbed orally, metabolised in the liver and excreted mainly in the urine.

Indications:

For relief of nonproductive cough due to minor throat and bronchial irritation, nasal and sinus congestion, sneezing and rhinorrhea associated with the common cold and allergic rhinitis.

Adverse Effects:

Drowsiness, lassitude, dizziness, nausea or vomiting, epigastric pain, blurred vision, confusion, difficulty in micturition, palpitations, headache, tinnitus, dryness of mouth, anorexia, and restlessness may occur occasionally.

Precautions:

- Caution in patients with asthma, hepatic function impairment, cardiovascular disease, glaucoma, hypertension, hyperthyroidism and prostatic hypertrophy.
- Check with physician if cough persists after medication has been used for 7 days.
- Caution when driving or doing jobs requiring alertness because of possible drowsiness.
- Safety for use in pregnancy has not been established.

Warning:

When used for treatment of cough and cold:

- Not to be used in children less than 2 years of age.
- To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

Contraindications:

- Use is not recommended for patients who are hypersensitive to other antihistamines or other sympathomimetics.
- It should not be used in nursing mothers.

Drug Interactions:

- Avoid alcohol or other central nervous system (CNS) depressants during therapy, unless prescribed or otherwise approved by physician.
- Concurrent use with monoamine oxidase (MAO) inhibitors is not recommended. This medication should not be administered during or within 14 days following the administration of a MAO inhibitor.
- Co-administration of atropine or other antimuscarinics may potentiate the muscarinic effects.

Overdosage:

Signs of overdose may include unsteadiness, flushing, shortness of breath, hallucinations, drowsiness, seizures, severe dryness of mouth, nose or throat, continuing headache and unusually slow or fast heartbeat, confusion, unusual excitement, nervousness, restlessness, or severe irritability.

Treatment consists of emesis or gastric lavage, and the necessary symptomatic and supportive therapy.

Dosage and Administration:

- | | | |
|----------|-----------------|---|
| Adults | - | Oral, 2 teaspoonfuls (10ml) three to four times a day as needed, not exceeding 40ml per day. |
| Children | - 2 to 5 years | - Oral, 1/2 teaspoonful (2.5ml) three to four times a day as needed, not exceeding 10ml per day |
| | - 6 to 12 years | - Oral, 1 teaspoonful (5ml) three to four times a day as needed, not exceeding 20ml per day. |

Storage : Keep container tightly closed. Store in cool, dry place below 30C. Protect from light. Keep out of reach of children.

Shelf life : 2 years

Pack size : Syrup x 90ml

Product registration holder: PRIME PHARMACEUTICAL SDN. BHD.

& Manufacturer 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah, 14000 Bukit Mertajam, Penang, Malaysia

MAL 19930042AZ

08/05/2023