



ASMIN EXPECTORANT

Description: Plain, cherry, flavoured, syrup.

Content

Each 5ml contains: Salbutamol Sulphate equivalent to Salbutamol 1.0mg
Guaiphenesin 50mg
Methylparaben 0.1% w/v
Propylparaben 0.01% w/v

Pharmacology:

Salbutamol is a direct-acting sympathomimetic amine. It acts selectively on beta-2-adrenergic receptors in the lungs to relax bronchial smooth muscle, thereby relieving bronchospasm and reducing airway resistance. Guaiphenesin has an expectorant action which reduces the viscosity of tenacious sputum, thereby facilitates the removal of mucus. It is readily absorbed orally, metabolised in the liver and excreted mainly in the urine.

Indications

For relief of productive and wheezing cough associated with bronchial asthma, bronchitis, pulmonary emphysema, bronchiectasis, and other obstructive pulmonary diseases.

Adverse Effects

Fast or pounding heartbeat, nausea, nervousness or restlessness, fine tremor of skeletal muscle (particularly the hands), palpitations, muscle cramps, headache and gastrointestinal discomfort may occur in some patients.

Warning & Precaution

- Caution in patients with coronary insufficiency, hypertension, diabetes mellitus, hyperthyroidism and phaeochromocytoma.
- Safety for use in pregnancy and lactation has not been established.
- Tocolysis: Serious adverse reactions including death have been reported after administration of terbutaline / salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

Contraindications:

Avoid in patients who are hypersensitive to sympathomimetics.

Drug Interactions

- Co-administration of glucocorticoid/adrenocorticoids may increase the metabolic clearance of adrenocorticoids; glucocorticoid dosage adjustment may be required.
- Concurrent use with adrenergic bronchodilators may result in mutual inhibition of therapeutic effects.
- Concurrent therapy with digitalis glycosides may increase the risk of cardiac arrhythmias.
- Concomitant administration of xanthines may increase CNS stimulation and other additive toxic effects.

Overdosage

Signs of overdose may include continuing or severe fast or pounding heartbeat, chest discomfort or pain, dizziness or light headedness, headache, increase in blood pressure, nausea or vomiting, trembling, and unusual anxiety, nervousness or restlessness.

Treatment consists of gastric lavage or administering a cardioselective beta-adrenergic blocker (with caution), if necessary for cardiac arrhythmias.

Dosage and Administration

Adults : Oral, 2 to 4 teaspoonfuls (10 to 20ml) two or three times a day, or as directed. The total daily dose should not exceed 130ml.
Children 2 to 5 years : Oral 1 to 2 teaspoonfuls (5 to 10ml) two or three times a day, or as directed.
6 to 12 years : Oral, 2 teaspoonfuls (10ml) two or three times a day, or as directed.

Storage Conditions : Keep container tightly closed. Store in a cool, dry place below 30°C. Protect from light.

Shelf life : 2 years

Presentation / Packing : Expectorant syrup X 90ml : 3.6L (EXPORT)

Manufactured & : PRIME PHARMACEUTICAL SDN BHD.

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

01 / 08 / 2018