

Artwork Document Information		Artwork Type [Please Tick /]		Sign / Stamp	
Product : T - DEXTROPHAN SYRUP 15MG/5ML SAP No. : Date : 29 April 2025		☐ Unit Box ☐ Outer Box ☐ Label ☒ Leaflet ☐ Fix-A-Form ☐ Shipper Carton ☐ Aluminium Foil ☐ Sachet ☐ Others : _____			
Document Category [Please Tick /]		Pharmacode / QR Code / Others Info			
☒ RA/NPRA Artwork ☐ Internal Artwork Circulation ☐ Others : _____ ☐ Recirculation Artwork ☐ Previous Approved Artwork ☐ Approval Copy Artwork ☐ Draft Artwork ☐ Clean Copy					
Remarks		Colour			
1. Font : Helvetica Regular (Spacing -8pt / Size -7pt) 2. Dimension : 161mm(w) x 200mm(h)		Black			
Colours shown on this proof are to indicate correct colour separations.					

DEXTROPHAN

Syrup 15mg/5ml

SOOTHES DRY, IRRITATING COUGH

DESCRIPTION

A brown, cherry flavoured syrup contains Dextromethorphan HBr 15 mg every 5 ml.
Preservative :
Methylparaben 0.21% w/v & Propylparaben 0.031% w/v.

INDICATION

Dextromethorphan is indicated for the symptomatic relief of nonproductive cough due to minor throat and bronchial irritation occurring with colds or inhaled irritants.

PHARMACODYNAMICS

Dextromethorphan is a cough suppressant which has a central action on the cough centre in the medulla. Although structurally related to morphine, dextromethorphan and other centrally acting cough suppressants have no analgesic properties and generally they have little sedative activity.

PHARMACOKINETICS

Dextromethorphan is well absorbed from the gastrointestinal tract. It is rapidly and extensively metabolised in the liver to dextrophan (active metabolite). It is also excreted primarily in the urine as unchanged dextromethorphan and demethylated metabolites including dextrophan. Dextrophan acts centrally to elevate the threshold for coughing. The onset of antitussive action occurs in 15-30 minutes after administration and the duration of action is up to 6 hours.

DOSAGE

Adults : 5 ml every 4 hours or 10ml every 6-8 hours. (max. 40ml/day)
Children 6 to 12 years : 2.5 ml every 4 hours or 5ml every 6-8 hours (max. 20ml/day)
Children 2 to 6 years: 1 - 2 ml every 4 hours or 2.5ml every 6-8 hours (max. 10ml/day).
Children under 2 years: Use is not recommended.

Route of administration: Oral

OVERDOSAGE

Dextromethorphan may produce central excitement and mental confusion. Toxic psychosis such as hyperactivity, visual and auditory hallucinations and respiratory depression have been reported to occur with very high doses. Dextromethorphan abuse and dependence may occur rarely especially following prolonged use of high dose. Induce emesis if patient is alert prior to 6 hours following ingestion. Precautions against aspiration must be taken, especially infants and small children. Gastric lavage may be carried out in removing unabsorbed drug. Respiratory depression should be treated promptly. The narcotic antagonist naloxone, is a specific antidote and should be administered intravenously if appropriate.

Up-to-date information about the treatment of overdose can be obtained from the National Poison Centre, University Sains Malaysia.

SIDE EFFECTS

Side effects with dextromethorphan appear to be rare and may include mild dizziness, drowsiness and gastrointestinal disturbances such as nausea, vomiting and stomach pain.

CONTRAINDICATIONS

Dextromethorphan is contraindicated in patients with hypersensitivity to the drug or any of the ingredients of the product. Risk-benefit should be considered in patients with asthma, productive cough and impaired hepatic function.

DRUG INTERACTIONS

Concurrent use with monoamine oxidase inhibitors (MAOI) including furazolidone and procarbazine may cause adrenergic crisis, collapse, coma, dizziness and excitation.
Concurrent use with alcohol and other central nervous system (CNS) depression producing medications may potentiate the CNS depressant effects of dextromethorphan.

PRECAUTIONS /WARNING

A persistent cough may be a sign of a serious condition. If cough persists after medication has been used for 7 days or if high fever, skin rash or continuing headache is present with cough, a physician should be consulted.
When used for treatment of cough and cold:
- Not to be used in children less than 2 years of age.
- To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

PREGNANCY & LACTATION

Dextromethorphan should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. It is not known whether dextromethorphan or its metabolites are excreted in breast milk.

PRESENTATION

90ml.

STORAGE CONDITIONS

Keep container tightly closed.
Store in a dry place below 30°C.
Protect from light.
Keep out of reach of children.
Jauhi daripada kanak-kanak

Shelf life : Please refer to outer package.

Product Registration Holder:

Duopharma Manufacturing (Bangi) Sdn. Bhd.
Lot No. 2, 4, 6, 8 & 10 Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

Duopharma Manufacturing (Bangi) Sdn. Bhd.
Lot 2 & 4, Jalan P/7, Section 13, Bangi Industrial
Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Revision Date: April 2025

