

|                       |                               |
|-----------------------|-------------------------------|
| Each 100 gm contains  |                               |
| Clobetasol Propionate | 0.05 gm in a white base cream |
| Methyl paraben        | 0.21 gm (as preservative)     |
| Propyl paraben        | 0.21 gm (as preservative)     |

### PHARMACODYNAMICS :

Clobetasol propionate decrease inflammation by stability leucocyte lysosomal membranes, preventing release of destructive acid hydrolases from leukocytes; inhibiting macrophage accumulation in inflamed area; reducing leukocyte adhesion to capillary endothelium; reducing capillary wall permeability and edema formation.

Precutaneous penetration of corticosteroid varies among individual patients and can be increased by the use of occlusive dressing, and by using different vehicles. Clobetasol and its metabolites are excreted in bile and in urine in animals.

Clobetasol propionate is a very active topical steroid which is of particular value in short courses for the treatment of the more resistant dermatoses eg psoriasis (excluding wide spread plaque psoriasis), recalcitrant eczemas, lichen planus, discoid lupus erythematosus and other conditions which do not respond satisfactorily to less active steroids.

Apply CLOBET CREAM to the affected area, once or twice daily until improvement occurs or as directed.

**PRECAUTIONS/ WARNINGS :**

- Long term continuous therapy with CLOBET CREAM in infants and children should be avoided since it may induce adrenal suppression.
- It should not be used extensively in pregnancy, either in large amounts or for prolonged periods.
- Appropriate chemotherapy should be used whenever secondary infections occurred.
- If used in childhood or on the face, courses should be limited if possible to 5 days and occlusion should not be used.
- The face, more than other areas of the body, may exhibit atrophic changes after prolonged treatment with potent topical corticosteroids. This must be borne in mind when treating such conditions as psoriasis, discoid lupus erythematosus and severe eczema.
- If applied to the eyelids, care is needed to ensure that the preparation does not enter the eye, as glaucoma might result.
- Topical corticosteroids may be hazardous in psoriasis for a number of reasons including rebound relapses, development of tolerance, risk of generalised pustular psoriasis and development of local or systemic toxicity due to impaired barrier function of the skin. If used in psoriasis, careful patient supervision is important.

As there are no adequate and well controlled studies in pregnant women, the safety of topical steroid during pregnancy has not yet been established. Topical steroid should be used only when the potential benefits outweighs the possible risk to the fetus.

The drug should not be used on extensive areas, in large amounts or for prolonged periods in pregnant women.

It is not known whether topical steroids are distributed into milk. Caution should be exercised when administering topical steroids to a nursing women.

Provided the weekly dosage is <50 g in adults, any pituitary-adrenal suppression is likely to be transient with a rapid return to normal values once the short course of steroid therapy has ceased. The same applies to children given proportionate dosage. Use of occlusive dressings increases the absorption of topical corticosteroids.

Prolonged and intensive treatment with highly-active corticosteroid preparations may cause atrophic changes, eg striae, thinning of the skin, and dilation of the superficial blood vessels, and particularly when occlusive dressings are used, or where skin folds are involved.

In some instances, treatment of psoriasis with corticosteroids (or its withdrawal) is thought to have provoked the pustular form of the disease. If signs of hypersensitivity appear, applications should be stopped immediately. As with other steroids, prolonged use of large amounts, or treatment of extensive areas can result in sufficient systemic absorption to produce the features of hypercorticism.

Rosacea, acne and perioral dermatitis, Primary cutaneous viral infections (eg. herpes simplex, chickenpox) and hypersensitivity to the preparation.

Because of topical application, the overdose would not occurred.

Store at temperature not exceeding 30°C. Keep medicine out of reach of children.

Aluminium tube 10 gm.

## October 2023



EER 001/D

|                                                        |                    |                            |
|--------------------------------------------------------|--------------------|----------------------------|
| Code : EER001/D                                        | Draft Revision : 5 | Final Artwork Revision : B |
| Name : EER001-D_Rev.B_Leaflet_Clobet cream_Malaysia.ai |                    |                            |
| Remark :                                               |                    |                            |

  

|           |               |                 |
|-----------|---------------|-----------------|
| Design by | Review by QCP | Approved by QCP |
| Signature | Signature     | Signature       |
| Date      | Date          | Date            |

  

|              |                |
|--------------|----------------|
| Review by RA | Approved by RA |
| Signature    | Signature      |
| Date         | Date           |

  

|               |                                                                                                                                     |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------|
| Specification | Size : 100 x 210 MM. (+/- 3 MM.)<br><br>Color : พิมพ์ 1 สี คือ สีดำ<br><br>Material : ใช้กระดาษปอนด์ น้ำหนัก 55 แกรม<br><br>Other : |
|---------------|-------------------------------------------------------------------------------------------------------------------------------------|