

YSP Antifungal Cream 20mg

Ingredient(s):

Each gram contains:

Miconazole Nitrate 20mg

Preservatives:

Methyl Paraben 1mg

Propyl Paraben 1mg

Action(s):

Pharmacodynamic

Miconazole Nitrate is a synthetic antifungal agent that is chemically unrelated to other fungicidal compounds. Although its pharmacologic mode of action has not been established, in vitro studies show that it possesses fungicidal action against *Candida albicans* and other species of this genus. It has also been found to inhibit the growth of common dermatophytes, *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Epidermophyton floccosum*, and the organism responsible for *Tinea versicolor*. Its therapeutic value appears to be due to its action primarily on the yeast cell membrane resulting in selective inhibition of the uptake of those components necessary for cell reproduction and growth. Clinical trials have shown it to be well-tolerated and effective in both pregnant and non-pregnant women as well as those taking oral contraceptives.

Pharmacokinetics

Miconazole is an imidazole antifungal agent. It has a wide antifungal spectrum and possesses some antibacterial activity. At concentration at 0.1 to 1 mg per ml, it inhibits the dermatophytes, *Epidermophyton floccosum*, *Microsporum canis*, *Trichophyton mentagrophytes* and *Trichophyton rubrum*, *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Paracoccidioides brasiliensis*, *Cryptococcus neoformans*, *Malassezia furfur* (*Pityrosporum orbiculares*), and some *Streptomyces* species. Concentrations of 1 to 10 mg per ml have been reported to inhibit other dermatophytes and *Candida*, and *Cladosporium*, and *Madurella* species, *Coccidioides immitis* and *Sporothrix schenckii*. Other sensitive fungi include *Phialophora pedrosoi* and *Aspergillus* and *Nocardia* species. Sensitive bacteria include *Staphylococci*, *Streptococci*, *Bacillus anthracis* and *Bacteroides fragilis*.

Following intravaginal administration of a single of dose of Miconazole (Cream or vaginal tablet) to healthy women, about 1% of the dose is recovered in urine and feces. There have no reports to data indicating that miconazole nitrate is absorbed following application to intact skin.

Indication(s):

For topical application in the treatment of *Tinea pedis* (athlete's foot), *Tinea cruris*, and *Tinea corporis* caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Epidermophyton floccosum*, and in the treatment of *Tinea versicolor*.

Dosage and Administration:

Sufficient Antifungal cream should be applied to cover affected areas twice daily (morning and evening) in patients with *Tinea pedis*, *Tinea cruris*, *Tinea corporis*, and cutaneous Candidiasis, and once daily in patients with *Tinea versicolor*.

Early relief of symptoms (2 to 3 days) is experienced by the majority of patients and clinical improvement may be seen fairly soon after treatment is begun; however, *Candida* infections and *Tinea cruris* and *corporis* should be treated for two weeks and *Tinea pedis* for one month in order to reduce the possibility of recurrence. If a patient shows no clinical improvement after a month of treatment, the diagnosis should be redetermined. Patients with *Tinea versicolor* usually exhibit clinical and mycological clearing after two weeks of treatment.

Mode of administration:

To be applied topically

Contraindication(s):

Patients known to be hypersensitive to this drug.

Precaution(s) / Warning(s):

1. Do not use in children under 2 years of age except under the advice and supervision of a doctor.
2. If irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist.
3. In patients on warfarin, caution should be exercised and anticoagulant effect should be monitored (see Interactions)

Please consult your physician/pharmacist before using this product if you are on the anticoagulant medicine warfarin, because bleeding from nose/gums or bruising may occur spontaneously

Interaction with other medicaments:

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and precautions)

Pregnancy and lactation:

Pregnancy

Miconazole nitrate has shown no teratogenic effects but is foetotoxic at high oral doses. Only small amounts of miconazole nitrate are absorbed following topical administration. However, as with other imidazoles, miconazole nitrate should be used with caution during pregnancy.

Lactation
Topically applied miconazole is minimally absorbed into the systemic circulation, and it is not known whether miconazole is excreted in human breast milk. Caution should be exercised when using topically applied miconazole products during lactation.

Side Effect(s) / Adverse Reaction(s):
Most complaints are vulvovaginal burning, itching or irritation. Other complaints such as vaginal burning, pelvic cramps, hives, skin rashes and headache occurred rarely. Irritation, burning, maceration and allergic contact dermatitis associated with application of Miconazole.

Symptoms and Treatment for Overdosage, and Antidote(s):
Overdosage in humans has not been reported to date.

Shelf-Life:
3 years from the date of manufacture.

Storage Condition(s):
Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture.

Packing(s):
Aluminium tube of 10g and 20g.
Plastic jar of 500g (for export only).
A white color cream.

YSP Antifungal Krim 20mg

Bahan Aktif:
Setiap gram mengandungi:
Miconazole Nitrate 20mg
Pengawet:
Methyl Paraben 1mg
Propyl Paraben 1mg

Indikasi:
Rawatan topikal untuk kurap kaki, kurap bahagian sulit, dan kurap badan yang disebabkan oleh *Trichophyton rubrum*, *Trichophyton mentagrophytes*, dan *Epidermophyton floccosum*, dan untuk rawatan panau.

Dos:
Untuk kurap kaki, kurap bahagian sulit, kurap badan, dan kandidiasis bahagian kulit, sapu krim yang mencukupi di bahagian yang terlibat dua kali sehari (pagi dan petang). Untuk rawatan panau, sapu krim yang mencukupi di bahagian yang terlibat satu kali sehari.

Kelegaan gejala (2 hingga 3 hari) akan dialami oleh majoriti pesakit dan tanda pemulihan mungkin dapat dilihat selepas rawatan dimulakan; Walau bagaimanapun, jangkitan *Candida*, kurap bahagian sulit dan kurap badan perlu dirawat selama dua minggu dan kurap kaki selama sebulan untuk mengurangkan risiko jangkitan berulang. Jika tiada tanda pemulihan secara klinikal selepas rawatan selama sebulan, diagnosis perlu ditentukan semula. Jangkitan panau biasanya akan menunjukkan tanda pemulihan secara klinikal dan mikologi selepas rawatan selama dua minggu.

無 黴 乳 膏

成份:
每公克中含:
Miconazole Nitrate 20mg

防腐劑:
Methyl Paraben 1mg
Propyl Paraben 1mg

適應症:
針對足癬 (香港腳)、股癬和體癬均有效,也有效於治療花斑癬 (汗斑)。

用法、用量:
足癬、股癬、體癬和皮膚念珠菌感染者每日兩次 (早晚) 塗於患處。花斑癬感染者每日一次塗於患處。

通常治療開始後的兩至三天會逐漸看到療效。念珠菌感染、股癬和體癬應治療兩週;足癬應治療一個月,以減少復發的可能性。如果患者在治療一個月後沒有得到改善,則應重新診斷。花斑癬通常在治療兩週後可以被清除。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679