



PRICOLAX TABLET (enteric-coated)

Description : Yellow, round, biconvex and enteric coated tablet, Diameter = 6.00 mm.

Content

Each tablet contains bisacodyl 5 mg

Indications

For constipation, pre & post operation evacuation and evacuation prior to radiology.

Pharmacology

Bisacodyl is a contact laxative which acts directly on the smooth intestinal musculature by stimulation of intramural nerve plexi. This increases peristalsis throughout the large intestine. It is poorly absorbed after oral administration. Up to 38% of the glucuronide conjugated bisacodyl has stated to be excreted in the urine in some patients. But most of the dose is excreted in the feces as the unconjugated hydrolysis product

Adverse Effects

May occasionally cause severe abdominal cramps, diarrhoea, belching or nausea.

Precaution

Avoid prolonged use as it may result in serious loss of water and electrolytes, muscular weakness and weight loss. Do not use products with potassium or magnesium salts of phosphates in patients with renal dysfunction.

No chewing or crushing tablets, swallowing whole because of enteric coating.

Not to be taken with milk or antacids.

Contraindications

Special precautions should be taken when used during pregnancy and in children.

Not recommended for children under 6 years old since patients in this group may have difficulty swallowing the tablet without chewing it.

Gastric irritation may occur if the enteric coating is destroyed by chewing.

Dosage and Administration

Adults and children above 6 years old : 1 - 2 tablets to be taken at bedtime.

Drink at least 6 to 8 full glasses (240ml each) of liquids every day when using laxative to aid stool softening and to protect against dehydration when large volume of water are lost with passage of stool.

To prepare patient for 'pre-post operation' and 'prior to radiology', 2 - 4 tablets to be taken the night before and one suppository to be inserted the following morning for complete evacuation of the intestine.

Storage : Keep container tightly closed.
Store in a cool, dry place, below 30°C.
Protect from light. Keep away from children.

Presentation/Packing : 100 x 10's / Alu-PVC blister
30's / bottle

Shelflife : 3 years

Manufactured & **PRIME PHARMACEUTICAL SDN.BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1,
Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam,
Penang, Malaysia.

MAL 19940044XZ

For further information, please consult your doctor or pharmacist.