

GERSKIN® TABLET

Gerskin tablets are used to anticipate and relieve flatulence or pain in the symptomatic management of gastric and duodenal ulcer and reflux esophagitis by neutralizing Hydrochloric Acid in the gastric secretion.

Ingredient(s):

Each tablet contains:

Magnesium Oxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	20mg

Action(s):

1. Magnesium Oxide and Aluminium Hydroxide Gel Dried can react chemically to neutralize or buffer existing quantities of stomach acid but has no direct effect on its output. This action results in increased pH value of stomach contents, thus providing relief of hyperacidity symptoms.
2. Magnesium Oxide and Aluminium Hydroxide Gel Dried can also reduce acid concentration within the lumen of the esophagus, thus causing an increase in intra-esophageal pH and a decrease in pepsin activity, which aids in the control of gastroesophageal reflux.
3. Dimethylpolysiloxane has antifoaming and water repellent properties. Its use in medicine as an antifatulent, is given specifically to promote coalescence of mucous-entrapped gas bubbles in the gastrointestinal tract and thereby facilitate their expulsion.
4. Small amount of the Gerskin tablets is absorbed from the intestine. The duration of action in fasting patients is 20 to 30 minutes. However, when the Gerskin tablets are given 1 hour after meals the acid-neutralizing effect may be prolonged up to 3 hours.
5. It is excreted in the renal and fecal: 15% to 30% of the salts formed are absorbed and excreted by kidney.

Indication(s):

For the treatment of hyperacidity, gastro-intestinal ulcer, gastritis, flatulence.

Contraindication(s):

Should not be administered to patients with renal failure, or who are severely debilitated.

Dosage and Administration:

Adults: 1 - 2 tablets, 3 - 4 times daily during meals or at bedtime.

Precaution(s) / Warning(s):

Do not take this product if you are presently taking a prescribed antibiotic containing any form of Tetracycline, Digoxin and Vitamin.

Side Effect(s) / Adverse Reaction(s):

Gerskin tablets are usually well tolerated, the side effects

which have been noted are rare, consisting of soft feces, diarrhea, nausea, vomiting, gastralgia and anorexia.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

5 years from the date of manufacture.

Product Description & Packing(s):

A round, concurred white tablet, impressed with a circle and

dot in the center on one side. “”

Plastic containers of 30's, 60's.

Plastic containers of 1000's, 1200's & 1500's.(For export only)

Blister packing of 10's x 10 and 10's x 100.

The information contained in this leaflet is limited.
Further information can be obtained from other sources
including your doctor or pharmacist.

GERSKIN® TABLET

Bahan Aktif:

Setiap tablet mengandungi:

Magnesium Oxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	20mg

Indikasi:

Untuk rawatan hiperasiditi, ulser gastrik, gastritis dan sebu perut.

Dos:

Dewasa: 1 - 2 tablet, 3 - 4 kali sehari semasa makan atau sebelum tidur.

克 酸 氣 錠

成份：

每錠含：

Magnesium Oxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	20mg

適應症：

胃、十二指腸潰瘍、胃酸過多、胃炎、鼓腸。

用法、用量：

成人每次 1~2 錠，每日 3~4 次，於進食中或就寢前服用。



Factory

Manufacturer and Marketing Authorization Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Office

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Tel: 03-89251215 (4 Lines) Fax: 03-89251298

No. 18, Jalan Wan Kadir, Taman Tun Dr. Ismail,
60000 Kuala Lumpur, Malaysia.
Tel: 03-77276390 (12 Lines)
Ordering line: 1 800 88 3027
Product info: 1 800 88 3679