

SHINOXOL[®] TABLET

Product Description:

A white colour round tablet, one side impressed with a score “

and the other side impressed with a mark “



Ambroxol, a metabolite of Bromhexine, is a mucolytic agent capable of increasing sputum volume and reducing its viscosity, improving breathing and reducing cough with no influence on gastric acid secretion.

Ingredient(s):

Each tablet contains:

Ambroxol Hydrochloride 30mg

Pharmacodynamics:

Ambroxol is a mucus-modifying drug, enhances both bronchial glandular cell secretions and the production of alveolar type II pneumocyte surfactant. It also stimulates ciliary motility and accelerates mucociliary transport.

Pharmacokinetics:

1. Shinoxol Tablet is a "mucus-modifier". Its active compound is Ambroxol. Several human studies showed that Ambroxol affects the secretomotor activity of serous glands, facilitates the repair of bronchial epithelium, and accelerates mucociliary transport and clearance.
2. Ambroxol, a metabolite of Bromhexine, has been found to stimulate the synthesis of lung surfactant, as the surfactant may play a role in airway mucus transport acting as a lubricant.
3. In several studies, Ambroxol also produced statistically significant symptomatic improvement, measured as difficulty in expectoration, coughing, presence of dyspnea and the auscultatory sign as compared to controls.
4. Absorption after oral administration of Ambroxol was found to be fast and complete. Elimination half-life in the blood was estimated as 20 - 25 hours in man. In man and rabbit, radioactive-labelled substance was excreted almost completely into the urine.

Indication(s):

Mucolytic agent used in the treatment of respiratory tract disorders associated with viscid mucus.

Dosage and Administration:

For adults: 1 tablet, thrice daily.

For long-term treatment, 1 tablet, twice daily.

To be taken with water after meals.

Route of administration:

Oral administration

Contraindication(s):

It is contraindicated in patients who have shown hypersensitivity to this drug.

Pregnancy and Lactation:

The benefits of the drug in pregnant women during first trimester should be weighed against a possible risk to the fetus.

Side Effect(s) / Adverse Reaction(s):

The side effects of Ambroxol include gastric discomfort, gastralgia, diarrhea, nausea, vomiting, constipation and dyspepsia. Other side effects such as skin rash, pruritus and erythema have also been reported.

Immune System Disorders Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Precaution(s) / Warning(s):

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms

appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdose, therefore, management of the patients should consist of symptomatic and supportive therapy.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Packing(s):

Plastic bottle of 200's and 1000's (for export only).
Blister packing of 10's x 10, 10's x 50 and 10's x 180.

SHINOXOL[®] TABLET

Bahan Aktif:

Setiap tablet mengandungi:
Ambroxol Hydrochloride 30mg

Indikasi:

Digunakan untuk merawat gangguan saluran pernafasan berhubung dengan mukus tebal.

Dos:

Untuk dewasa: 1 tablet 3 kali sehari.
Rawatan jangka panjang: 1 tablet 2 kali sehari.

暢 痰 錠

成份：

每錠含：
Ambroxol Hydrochloride 30mg

適應症：

化痰。

用法、用量：

成人每次1錠，每日3次。
長期治療：每次1錠，每日2次。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679