

## PROPOSED INSERT

hovid

### ACTIHIST-CO SYRUP

VIACtxx-PC (MY)

#### DESCRIPTION

Red, transparent liquid with tutti-frutti flavour.

#### COMPOSITION

Each 5 ml contains:

|                               |          |
|-------------------------------|----------|
| Tripolidine Hydrochloride     | 1.25 mg  |
| Pseudoephedrine Hydrochloride | 30.00 mg |
| Dextromethorphan Hydrobromide | 10.00 mg |

#### PHARMACODYNAMICS

##### Pseudoephedrine:

Pseudoephedrine is a sympathomimetic agent. It is used as a bronchodilator and peripheral vasoconstrictor in preparations for the relief of nasal and bronchial congestion. It has a stimulant action on the respiratory center. It acts on alpha-adrenergic receptors in the mucosa of the respiratory tract.

##### Tripolidine:

Tripolidine is an antihistamine. It is an antagonist of the H-1 receptor effects of histamine. It acts by competing with histamine for H-1 receptor sites on effector cells. It diminishes or abolishes most of the actions of histamine. However, it does not antagonise its stimulant action on gastric secretion which is mediated by H-2 receptors. The action of histamine most readily affected is that on the involuntary muscle of the intestines, bladder, uterus and bronchioles.

The antimuscarinic actions of Tripolidine provide a drying effects on the nasal mucosa.

##### Dextromethorphan:

Dextromethorphan Hydrobromide is a cough suppressant. It exerts its effects by acting on the cough center in the medulla.

#### PHARMACOKINETICS

##### Absorption:

Well absorbed from the gastrointestinal tract.

##### Metabolic Reactions:

Undergoes hepatic metabolism

##### Excretion:

Undergoes renal excretion. Pseudoephedrine is largely excreted unchanged in the urine together with small amounts of its hepatic metabolite while Tripolidine is mainly excreted as metabolites in the urine. Both Pseudoephedrine and Tripolidine are also excreted in breast milk.

#### INDICATIONS

This medication is indicated for the symptomatic treatment of unproductive cough and nasal congestion associated with allergic rhinitis, vasomotor rhinitis, hay fever and common cold.

#### CONTRAINDICATIONS

This medication should not be given to premature infants or neonates, patients undergoing anaesthesia with cyclopropane, halothane or other halogenated anaesthetics and patients with hyperthyroidism, cardiovascular disease or acute attacks of asthma. It should not be used concurrently with monoamine oxidase inhibitors.

#### PRECAUTIONS

- This medication should be used with caution in hypersensitive patients and in geriatrics. It should be carefully considered when the following medical problems exist: Diabetes mellitus, close-angle glaucoma, hypertension, peptic ulcer and renal function impairment.
- Safety for use in nursing mothers has not been established.
- Not to be used in children less than 2 years of age.
- To be used with caution and doctor's advice in children 2 to 6 years of age.

#### PREGNANCY AND LACTATION

Not recommended in pregnancy and lactation.

#### MAIN SIDE/ADVERSE EFFECTS

Drowsiness, insomnia, urinary retention, nausea, vomiting and allergic reactions.

#### DRUG INTERACTIONS

Concurrent use with the following drugs is not recommended: Beta-adrenergic blocking agents, digitalis glycosides, guanethidine, methyl dopa, alcohol, CNS depressants, tricyclic antidepressants, antimuscarinics and ototoxic medications.

#### OVERDOSAGE

##### Clinical features:

Headache, nausea, vomiting, dryness of the mouth, tachycardia, respiratory depression, coma.

##### Treatment:

Emesis or gastric lavage, followed by activated charcoal. Supportive and symptomatic measures. Adrenaline and physostigmine are contraindicated.

#### DOSAGE AND ADMINISTRATION

##### Adults:

Oral, 10 ml three times a day as needed.

##### Paediatrics:

2 to 6 years of age: Oral, 2.5 ml three times a day.

6 to 12 years of age: Oral, 5 ml three times a day as needed.

Note: Geriatric patients may be more sensitive to the effects of the usual adult dose.

This medication may be taken with food, water or milk to minimize stomach irritation.

The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 25°C. Protect from light.

##### Presentation/Packing:

Syrup x 60 ml, 100 ml, 120 ml,  
3.6 Litres (For export only).

Product Registration Holder /  
Manufactured by: HOVID Bhd.  
121, Jalan Tunku Abdul Rahman,  
30010 Ipoh, Malaysia.

hovid