
MICO CREAM

Name and Strength of Active Ingredient (s):

Miconazole Nitrate BP 2.0 % w/w

Preservatives:

Methyl Paraben 0.15% w/w

Propyl Paraben 0.08% w/w

Product Description:

White colour cream.

Pharmacodynamics:

Pharmacotherapeutic classification: (Antifungals for dermatological/topical use; imidazole derivative) ATC code: D01AC02.

Miconazole is an imidazole antifungal agent and may act by interfering with the permeability of the fungal cell membrane. It possesses a wide antifungal spectrum and has some antibacterial activity.

Pharmacokinetics:

Absorption: There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically.

Distribution: Absorbed miconazole is bound to plasma proteins (88.2%) and red blood cells (10.6%).

Metabolism and Excretion: The small amount of miconazole that is absorbed is eliminated predominantly in faeces as both unchanged drug and metabolites.

Indications:

Mico Cream is indicated for the topical treatment of the skin infections due to susceptible fungi (dermatophytes and yeasts) such as tinea pedis (athlete's foot), tinea cruris, tinea corporis (ringworm), tinea versicolor (white spots), and candidiasis.

Recommended Dosage:

Apply a thin layer of Mico Cream sufficient to cover the affected areas and rub well into the skin until the cream has been completely absorbed. The cream should be applied once or twice a day, in the morning and evening, and continued without interruption until symptoms have been completely disappeared (usually 2 to 5 weeks, depending on the infecting organism and the site of infection).

Mode of Administration:

Topical.

Contraindications:

Mico Cream is contraindicated in individuals with a known hypersensitivity to miconazole or another ingredient in this product.

Warning and precautions:

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see interactions).

Discontinue treatment if sensitisation or irritation occurs during use.

DO NOT USE IN THE EYES.

Do not use in children below 2 years old except under the advice and supervision of a doctor.

If there is no improvement, discontinue and consult doctor or pharmacist.

- Within 4 weeks for athletes' foot

- Within 2 weeks for jock itch.

Interactions with other Medicaments

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and anticoagulant effect should be monitored (See Warnings and Special Precautions).

Pregnancy and Precaution:*Pregnancy:*

In animals, miconazole nitrate has shown no teratogenic effects but is foetotoxic at high oral doses. Only small amounts of miconazole nitrate are absorbed following topical administration. However, as with other imidazoles, miconazole nitrate should be used with caution during pregnancy.

Lactation:

Topically applied miconazole is minimally absorbed into the systemic circulation, and it is not known whether miconazole is excreted in human breast milk. Caution should be exercised when using topically applied miconazole products during lactation.

Adverse Effects:

Mico Cream is usually well tolerated. However, local irritation and sensitivity reactions may occasionally occur.

Overdose and Treatment:*Symptoms*

Cutaneous use: Excessive use can result in skin irritation, which usually disappears after discontinuation of therapy.

Treatment

Mico Cream is intended for cutaneous use, not for oral use. If accidental ingestion of large quantities of the product occurs, use appropriate care.

Effects on Ability to Drive and Use Machines:

Not Applicable.

Storage Conditions:

Store below 30°C. Protect from freezing, heat and direct light.

Packaging Available:

15 gm tube, 15 gm x 25 tubes and 15 gm x 50 tubes.
20gm x 10 tubs.

MAL Number:

MAL19940219AZ

Manufactured by & Product Registration Holder:

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

17 October 2023