

Somin Tablet 2mg

Ingredient(s):

Each tablet contains:

Dexchlorpheniramine Maleate 2mg

Pharmacodynamics:

1. Somin is an antihistamine with anticholinergic properties. It is capable to compete with histamine for receptor sites on effector cells and is of value clinically in the prevention and relief of many allergic manifestations.
2. In vitro and in vivo assays of the antihistamine potencies of the optically active isomers of Chlorpheniramine demonstrate that the predominant activity is in the dextro-isomer. The dextro-isomer is approximately 2 times more active than the racemic compound.

Pharmacokinetics

1. Dexchlorpheniramine is well absorbed after oral administration. It is mainly metabolised in the liver and some in the renal. Half-life is 12 to 15 hours.

Indication(s):

For symptomatic relief of allergic condition, for example treatment of perennial and seasonal allergic rhinitis, vasomotor rhinitis, allergic conjunctivitis, mild or uncomplicated allergic skin manifestation of urticaria and angioedema, amelioration of allergic reaction to blood or plasma, dermographism. Somin is also indicated as therapy for anaphylactic reaction adjunctive to Epinephrine and other standard measure after the acute manifestations have been controlled.

Dosage and Administration:

Adults: 2mg 4 - 6 hourly as required.

Children:

2 to 5 years: 0.5mg every 4 to 6 hours up to a maximum of 3mg in 24 hours.

6 to 11 years: 1mg every 4 to 6 hours up to a maximum of 6mg in 24 hours.

Contraindication(s):

Somin tablet is contraindicated in patients during acute attacks of asthma and premature infants or neonates.

Route of Administration:

Oral

Precaution(s) / Warning(s):

Because of Dexchlorpheniramine Maleate having antimuscarinic properties, it should be used with care in conditions such as narrow angle glaucoma, urinary retention, and prostatic hypertrophy. MAOIs may enhance the antimuscarinic effects of antihistamines. Because it may cause drowsiness, patients so affected should not drive or operate machinery. Dexchlorpheniramine Maleate has an atropine-like action and therefore products containing it should be used with caution in patients with a history of bronchial asthma, increased intraocular pressure, hyper-thyroidism, cardiovascular disease, and hypertension.

Drug Interaction(s):

Somin prolongs and intensifies Monoamine Oxidase (MAO) inhibitors effect which may lead to severe hypotension. Concomitant use of Somin with alcohol, tricyclic antidepressants, barbiturates or other CNS depressants may potentiate the sedative effect.

Side Effect(s) / Adverse Reaction(s):

The most common side effect is sedation, varying from slight drowsiness to deep sleep, and including lassitude, dizziness, and incoordination. Other side effects include gastrointestinal disturbances such as nausea, vomiting, diarrhea or constipation, anorexia or increased appetite, and epigastric pain.

Symptoms and Treatment for Overdosage, and Antidote(s):

In the treatment of overdosage, emesis should be induced if vomiting has not occurred spontaneously, Syrup of Ipecac is recommended. Precautions against aspiration must be taken, especially in infants and children. Vasopressors may be used to treat hypotension (Epinephrine should not be used since it may further lower the blood pressure); other measures include oxygen and intravenous fluids.

Shelf-Life:

Plastic bottle : 5 years from the date of manufacture.

Blister packing : 3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product description:

A pink color round tablet, one side is impressed with a "C" and the other side with a score.



Packing(s):
Plastic bottle of 1000's and 2000's (for export only)
Blister packing of 10's x 10 and 10's x 100

Somin Tablet 2mg

Bahan Aktif:
Setiap tablet mengandungi:
Dexchlorpheniramine Maleate 2mg

Indikasi:
Memberi kelegaan simptomatik untuk keadaan alahan, misalnya untuk rawatan rinitis allergik perenial dan seasonal, rinitis vasomotor, konjunktivitis allergik, alahan kulit yang ringan dengan manifestasi urticaria dan angioedema, pelegaan tindakbalas alahan terhadap sediaan darah atau plasma serta dermagrafisma. Juga diberi sebagai terapi untuk tindakbalas anafilaktik di samping pemberian Epinephrine dan rawatan sokongan lain selepas keadaan akut sudah terkawal.

Dos:
Dewasa: 2mg setiap 4 – 6 jam mengikut keperluan.
Kanak-kanak:
2 -5 tahun: 0.5mg setiap 4 – 6 jam sehingga maksima 3mg dalam 24 jam.
6 -11 tahun: 1mg setiap 4 – 6 jam sehingga maksima 6mg dalam 24 jam.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679