

MACGEL[®] TABLET

Ingredient(s):

Each tablet contains:

Magnesium Hydroxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	25mg

Pharmacodynamics

- Macgel Tablets are used to anticipate and relieve flatulence or pain in the symptomatic management of gastric and duodenal ulcer and reflux esophagitis by neutralizing hydrochloric acid in the gastric secretion.
- Magnesium Hydroxide and Aluminium Hydroxide Gel Dried can react chemically to neutralize or buffer existing quantities of stomach acid but has no direct effect on its output. This action results in increase pH value of stomach content, thus providing relief of hyperacidity symptoms. Magnesium Hydroxide and Aluminium Hydroxide Gel Dried can also reduce acid concentration within the lumen of the esophagus, thus causing an increase in intra-esophageal pH and a decrease in pepsin activity, which aids in the control of gastro-esophageal reflux.
- Dimethylpolysiloxane has antifoaming and water repellent properties. Its use in medicine as an antifatulent, is given specifically to promote coalescence of mucous-entrapped gas bubbles in the gastrointestinal tract and thereby facilitate their expulsion.
- Small amount of the Macgel tablets is absorbed from the intestine. The duration of action in fasting patients is 20 to 30 minutes. However, when Macgel Tablets are given 1 hour after meals the acid-neutralizing effect may be prolonged up to 3 hours.
- It is excreted in the renal and fecal; 15% to 30% of the salts formed are absorbed and excreted by kidney.

Pharmacokinetics

Small amount of Macgel Tablet are absorbed from the intestine. The duration of action in fasting patients is 20 ~ 30 minutes. However, when the Macgel tablet are given 1 hour after meals the acid-neutralizing effect may be prolonged up to 3 hours. It is excreted in the renal and fecal; 15 to 30% of the salts formed are absorbed and excreted by kidney.

Indication(s):

Hyperacidity, peptic and duodenal ulcer, gastritis, flatulence.

Route of administration: Oral

Dosage and Administration:

Adults: 2-4 tablets, 4 times daily after meals and at bedtime. Chew tablets before swallowing or swallow whole.

Contraindication(s):

Macgel is contraindicated in patients who have shown hypersensitivity to the drug.

Precaution(s) / Warning(s):

Macgel should not be given in conjunction with tetracycline antibiotics.

Side Effect(s) / Adverse Reaction(s):

Macgel tablets are usually well tolerated. The side effects which have been noted are rare, consisting of soft feces, diarrhea, nausea, vomiting, gastralgia and anorexia.

Symptoms and Treatment for Overdosage, and Antidote(s):

In the event of accidental overdosage, vomiting and purging should be induced.

Storage Condition(s):

Store at temperature below 30°C.

Protect from light and moisture.

Shelf-Life:


3 years from the date of manufacture.

Packing(s):

Blister packing of 10's x 2, 10's x 10, 10's x 100 and 10's x 120.

Product description:

A round tablet, white color with “YS” mark at one side and pink color with a score at the other side “



The information contained in this leaflet is limited. Further information can be obtained from other sources including your doctor or pharmacist.

MACGEL[®] TABLET

Bahan Aktif:

Setiap tablet mengandungi:

Magnesium Hydroxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	25mg

Indikasi:

Hiperasiditi, ulser peptik dan duodenal, gastritis dan sebu perut.

Dos:

Dewasa: 2 - 4 tablet, 4 kali sehari, selepas makan dan sebelum tidur.
Kunyahkan tablet sebelum telan atau telan seluruhnya.

皆 樂 錠

成份:

每錠含:

Magnesium Hydroxide	200mg
Aluminium Hydroxide Gel Dried	200mg
Dimethylpolysiloxane	25mg

適應症:

胃酸過多、胃、十二指腸潰瘍、胃炎、腹脹。

用法 . 用量:

成人每日4次，每次2~4錠，於飯後及睡前咀嚼吞服。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679