

BROZIL[®] Capsule 300mg

Brozil Capsule contains Gemfibrozil, a synthetic antihyperlipidemic agent which is structurally related to Clofibrate. It has been shown to be 10 times as potent in reducing triglycerides in animal studies and three times more active than Clofibrate in reducing platelet aggregation. Preliminary studies in human volunteers indicate that Gemfibrozil is a safe and effective antihyperlipidemic agent in selected cases of primary hyperlipidemia.

Ingredient(s):

Each capsule contains:

Gemfibrozil 300mg

Coloring agents for capsule:

Brilliant Blue FCF

Cochineal Red A

Pharmacodynamics:

- a. Gemfibrozil reduces plasma triglycerides (very low-density lipoprotein, VLDL) concentration and increases high-density lipoprotein (HDL). The mechanism of this action is not completely understood but may involve inhibition of peripheral lipolysis; reduced hepatic extraction of free fatty acids which reduces hepatic triglycerides production; inhibition of synthesis and increased of clearance of VLDL carrier, apolipoprotein B, which also reduces VLDL production; and, according to animal studies, reduced incorporation of long chain fatty acids into newly formed triglycerides, accelerated turn-over and removal of cholesterol from the liver (stimulated incorporation of cholesterol precursors into liver sterols), and increased extraction of cholesterol in the feces.
- b. Gemfibrozil is a lipid regulating agent which lowers elevated serum lipids primarily by decreasing serum triglyceride with a variable reduction in total serum cholesterol. These decreases occur primarily in a very low density lipoprotein (VLDL) fraction. It has been reported that Gemfibrozil may increase levels of high density lipoprotein (HDL) subfractions, HDL² and HDL³, as well as apolipoprotein AI and AII.

Pharmacokinetics:

- a. Decreases in LDL are also observed although the magnitude of this change depends upon the type of hyperlipoproteinemia being treated. In the US multicenter study in which over 100 patients received Gemfibrozil 1200 mg daily for 2 years, serum total cholesterol decreases by 19%, 18% and 12% in type IIa, IIb, and IV patients, respectively; corresponding decreases in triglycerides being 34%, 43% and 39%; decreases in LDL cholesterol being 23%, 20%, and 11% whereas increases in HDL cholesterol averaged 0%, 11%, and 15%.
- b. Gemfibrozil is a well absorbed in the gastrointestinal tract after oral administration. Peak plasma level occur in 1 ~ 2 hours with a plasma half-life of 1.5 hours following single doses, and 1.3 hours following multiple doses. Plasma level appear proportional to dose and do not demonstrate accumulation over time following multiple doses.
- c. Gemfibrozil mainly undergoes oxidation of the ring methyl group to successively form a hydroxymethyl and carboxyl metabolite. Approximately 70% of the human dose is excreted in the urine, mostly of the glucuronide conjugate, with less than 2% excreted as unchanged Gemfibrozil. Six percent of the dose is accounted for in the feces.
- d. Unlike, Clofibrate, Gemfibrozil has no effects on plasma fibrinogen, nor does it influence glucose tolerance.

Indication(s):

Hyperlipidemia.

Dosage and Administration:

2 capsules twice daily, preferably 30 minutes before the morning and evening meals.

To be dispensed on physician's prescription.

Route of Administration: Oral

Contraindication(s):

1. Hypersensitivity to Gemfibrozil.
2. Alcoholism.
3. Pre-existing gallstones.
4. Primary biliary cirrhosis.

Precaution(s) / Warning(s):

1. Before instituting Brozil Capsule therapy, every attempt should be made to control serum lipids with appropriate diet, exercise, weight loss in obese patients, and to control other medical problems such as diabetes mellitus and hypothyroidism.
2. Pre-treatment laboratory studies should be performed to ensure that patients have abnormal levels of serum lipids. Periodic determination of serum levels should be obtained during administration. The drug should be withdrawn after 3 months if the lipid response is inadequate.
3. Brozil Capsule should be used with caution in patients with impaired hepatic function.
4. Safety and efficacy in children have not been established.
5. Concurrent use of anticoagulants with Gemfibrozil may significantly increase the anticoagulant effect of these medications; adjustments of anticoagulant dosage based on frequent prothrombin-time determinations are recommended.
6. Concurrent use of Lovastatin with Gemfibrozil may be associated with an increased risk of rhabdomyolysis, significant increases in creatine kinase concentrations, and myoglobinuria that leads to acute renal failure; may be seen as early as 3 weeks or as late as several months after initiation of combined therapy; monitoring of creatine kinase has not been shown to prevent severe myopathy or renal damage.
7. It should be used with caution in patients with impaired severe renal function. Reduced clearance leads to increased incidence of side effects; reduced dosage is recommended.

Pregnancy and Lactation:

1. Mothers taking the drug should not breast-feed.
2. It should be used only if potential benefit justifies the potential risk to the fetus.

Symptoms and Treatment of Overdose:

While there has been no reported cases of overdosage, symptomatic measures should be taken if it occurs.

Side Effect(s) / Adverse Reaction(s):

Gemfibrozil is well tolerated in most patients. The most frequent side effects noted in clinical trials consist of changes in bowel function, including abdominal pain, diarrhea, and occasional nausea in 3-5% of patients. Less common side effects include muscle tenderness and skin rashes. Biochemical changes are uncommon, that include eosinophilia, decreases in the plasma of alkaline phosphatase and occasionally rises in transaminases. Gemfibrozil potentiates the effects of oral anticoagulants, and may increase biliary lithogenicity with a predicted long-term increase in the incidence of gallstones.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Packing(s):

Blister packing of 10's x 10, 10's x 50, 10's x 60 and 10's x 100.

Product Description:

A size #0 maroon cap and white body hard gelatin capsule, the cap and body are marked with "G300"

**Y. S. P.**

Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679