

Opella Healthcare Italy

PACKAGING TEAM

Code:

936892

Update:

V02 - 11-04-2025

GMID code:

715833

Current item code:

845856

Product/Item type:

Istruzione

Maalox PET

250 ml

Country:

Malaysia

Artwork by:

Origgio

Plant:

Origgio

Technical Data

Format:

Stesa 157 x 315 mm

Plant barcode:

1047

Number of Colours:

1

K

Fonts:

Helvetica Regular / Bold

Minimum point size of text:

8 pt

Layout of Cutting:

Packaging Line:

L 010

GDO - Graphic Department Origgio

Opella Healthcare Italy S.r.l. - Viale Europa, 11

21040 Origgio (VA) - Italy - Tel. +39 - 02 96 10 559

Impianti di proprietà Opella Healthcare Italy S.r.l.

Vietata la manomissione - Rendere dopo la stampa

Maalox®Plus Suspension



Composition:
To provide symptomatic relief of hyperacidity plus alleviation of gas symptoms, each teaspoonful (5ml) contains:

Active ingredients	Per teaspoonful (5ml)
Magnesium hydroxide	200mg
Dried aluminium hydroxide gel (equiv. to aluminium hydroxide)	230mg (175mg)
Simethicone	25 mg

Pharmacological properties:
Pharmacodynamic properties
Pharmacotherapeutic group: Antacids with antifatulents; ATC code: A02AF02.

Mechanism of action
The antacid effect takes place in the stomach where, at the appropriate dose, magnesium hydroxide and aluminum hydroxide neutralize gastric acid production. The constipating effect of aluminum hydroxide is counteracted by the laxative effect of magnesium hydroxide. Simeticone has an antifatulent effect.

Pharmacokinetic properties

Absorption
When administered orally, aluminum hydroxide and magnesium hydroxide react with the hydrochloric acid produced in the stomach to form salts, which are only partly absorbed.

Excretion
Absorbed salts are excreted mainly in the urine.

Indications:
As an antacid for symptomatic relief of hyperacidity associated with the diagnosis of peptic ulcer, gastritis, peptic esophagitis, gastric hyperacidity, heartburn, or hialal hernia. As an antifatulent to alleviate the symptoms of gas, including postoperative gas pain.

Directions for Use:
2-4 teaspoonfuls, 4 times a day, taken 20 minutes to 1-hour after meal and at bedtime, or as directed by a physician. Maalox® Plus are not recommended for children. Route of administration: Oral

Adverse Effects:
The frequency of the undesirable effects listed below is defined using the following conventions:
Common (≥1/100 to <1/10); uncommon (≥1/1 000 to <1/100); rare (≥1/10 000 to <1/1 000); very rare (<1/10 000); not known (cannot be estimated from the available data).

Immune system disorders:
Frequency not known: angioedema, anaphylactic reactions, hypersensitivity reactions, urticaria, pruritus.

Gastrointestinal disorders:
Uncommon: diarrhea or constipation.
Frequency not known: abdominal pain.

Metabolism and nutrition disorders:
Very rare: hypermagnesemia, which has been observed with prolonged use in patients with kidney impairment. Frequency not known: hyperaluminemia, hypophosphatemia, in prolonged use or at high doses, or even normal doses of the medicinal product in patients on low-phosphorus diets, or in infants (0 to 24 months), which may result in increased bone resorption, hypercalciuria or osteomalacia.

Patient Warnings:
Do not take more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than two weeks or use if you have kidney disease except under the advice and supervision of a physician. Aluminum hydroxide may cause constipation and magnesium salts overdose may cause hypomotility of the intestine; large doses of this medicinal product may trigger or aggravate intestinal obstruction and ileus in patients with higher risk, such as those with renal impairment, primary constipation, impaired intestinal motility, infants (0 to 24 months) and the elderly. Aluminum hydroxide is not well absorbed from the gastrointestinal tract, and systemic effects are therefore rare in patients with normal renal function. However, excessive doses or long-term use, or even normal doses in patients with low-phosphorus diets or in infants (0 to 24 months), may lead to phosphate depletion (due to aluminum-phosphate binding) accompanied by increased bone resorption and hypercalciuria with the risk of osteomalacia. Medical advice is recommended in case of long-term use or in patients at risk of hypophosphatemia. In patients with renal impairment, plasma levels of aluminum and magnesium tend to increase, which causes hyperaluminemia and hypermagnesemia respectively. In these patients, long-term exposure to high doses of aluminum and magnesium salts may lead to encephalopathy, dementia, microcytic anemia or it may worsen dialysis-induced osteomalacia. Medical monitoring is required in patients with mild to moderate renal impairment, and long-term use of the medicinal product should be avoided. Administration of this medicinal product is contraindicated in patients with severe forms of renal failure. Aluminum hydroxide may be unsafe for patients with porphyria who are undergoing hemodialysis; Maalox® Plus is contraindicated in patients with porphyria. Due to its composition, Maalox® Plus does not generally affect bowel function. However, in some particularly sensitive patients and at high dosages, accelerated intestinal transit may occur.

Opella Healthcare Italy

PACKAGING TEAM

Code:

936892

Update:

V02 - 11-04-2025

GMID code:

715833

Current item code:

845856

Product/Item type:

Istruzione

Maalox PET

250 ml

Country:

Malaysia

Artwork by:

Origgio

Plant:

Origgio

Technical Data

Format:

Stesa 157 x 315 mm

Plant barcode:

1047

Number of Colours:

1

K

Fonts:

Helvetica Regular / Bold

Minimum point size of text:

8 pt

Layout of Cutting:

Packaging Line:

L 010

GDO - Graphic Department Origgio

Opella Healthcare Italy S.r.l. - Viale Europa, 11

21040 Origgio (VA) - Italy - Tel. +39 - 02 96 10 559

For Opella use only

Impianti di proprietà Opella Healthcare Italy S.r.l.

Vietata la manomissione - Rendere dopo la stampa

This medicinal product contains 448 mg sorbitol in 10 mL (2 teaspoons). Patients with hereditary fructose intolerance (HFI) should not take/be given this medicinal product.
This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.
Pediatric populations
Use of magnesium hydroxide in small children, particularly those with kidney impairment or dehydration, may result in hypermagnesemia.

- Contraindications:**
- Hypersensitivity to the active substances or to any of the excipients.
 - Patients with porphyria.
 - Severe renal insufficiency (creatinine clearance less than 30 mL/min).
 - Contraindicated in children.
 - Cachexia.

Drug Interaction:
Because Al and Mg salts reduce the gastrointestinal absorption of tetracyclines, the use of Maalox® Plus should be avoided during oral tetracyclines therapy.
The use of antacids containing aluminium can reduce the absorption of drugs, in particular H₂ antagonists, atenolol, cefdinir, cefpodoxime, chloroquine, tetracyclines, dasatinib monohydrate, diflunisal, digoxin, dexamethasone, bisphosphonates, eltrombopag olamine, elvitegravir, ethambutol, fluoroquinolones, glucocorticoids, indomethacin, iron salts, isoniazide, ketoconazole, levothyroxine, lincosamides, metoprolol, nilotinib, phenothiazine neuroleptics, penicillamines, propranolol, raltegravir potassium, rilpivirine, riociguat, rosuvastatin sodium fluoride, and antiviral treatments in combination with tenofovir alafenamide fumarate/emtricitabine/sodium bictegrovir.
As magnesium hydroxide administration results in urine alkalization, increased excretion of salicylates has been observed in the event of concomitant use.
Caution is advised when this medicinal product is used concomitantly with polystyrene sulfonate (Kayexalate) due to the potential risk of reduced effectiveness of the resin in binding potassium, of metabolic alkalosis in patients with renal impairment (reported with aluminum hydroxide and magnesium hydroxide) and of intestinal obstruction (reported with aluminum hydroxide).
Aluminum hydroxide and citrates may cause hyperaluminemia, especially in patients with renal impairment.
As a precaution, allow at least 2 hours (4 for fluoroquinolones) to elapse between taking any oral drug and Maalox® Plus.
Concomitant use with quinidine may increase the serum levels of quinidine and lead to quinidine overdose.
Concomitant use of aluminium hydroxide and citrates may result in an increased aluminum levels, especially in patients with renal impairment.
Urine alkalization following magnesium hydroxide administration may affect the excretion of some drugs; increased excretion of salicylates has been observed.

Pregnancy and Lactation:
There are no available data on Maalox® Plus use in pregnant women. No conclusions can be drawn regarding whether or not Maalox® Plus is safe for use during pregnancy.
Pregnancy
The medicinal product should only be used if absolutely necessary, under direct medical supervision, after having evaluated the expected benefits for the mother compared with the possible risk to the fetus or infant.
Breast-feeding
Because of the limited maternal absorption when taken in accordance with the indicated dosage regimen, aluminum hydroxide and its magnesium salt combinations are considered compatible with breast-feeding.

Effects on ability to drive and use machines:
Maalox® Plus has no influence on the ability to drive and use machines.

Overdose:
There is very limited experience with deliberate overdose. Cases of overdose with aluminum salts are more likely to occur in patients with severe chronic renal impairment, with the following symptoms: encephalopathy, seizures and dementia, hypermagnesemia.
The most frequently reported symptoms of acute overdose with aluminum hydroxide and in combination with magnesium salts include diarrhea, abdominal pain and vomiting.
Large doses of this medicinal product may cause or aggravate intestinal obstruction and ileus in patients at risk.
Treatment should be symptomatic and include general supportive measures.
Aluminum and magnesium are eliminated through the urinary route; treatment of magnesium overdose consists of rehydration and forced diuresis. In case of renal function deficiency, hemodialysis or peritoneal dialysis is necessary.

Description:
White suspension with odour and taste of lemon.

Supplied:
Maalox® Plus Suspension is available in bottles of 180ml, 250ml, 355ml, 500ml.
Not all presentations are marketed.

Storage condition:
Store below 30°C

Shelf-life:
Do not utilize after the expiry date indicated on the packaging.

Keep this and all drugs out of the reach of children.
Contain domiphen bromide 0.42mg/5ml as preservative.

Contains approximately 0.174 mg sodium per teaspoonful. To be taken into consideration by patients on a controlled sodium diet.

Manufactured by:
Opella Healthcare Italy S.r.l.
Viale Europa, 11
21040 Origgio (Va) - ITALY
Date of revision: April 2025
(based on IT SmPC June 2022)