

DRUG INTERACTIONS:

Not Applicable

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION:

Adult: 1 to 2 tablets to be chewed as required, after or between meals and at bedtime; or as directed by the physician.

Children: Over 6 years; ½ adult dose

Below 6 years: Not recommended otherwise as consulted by the physician.

SYMPTOMS AND TREATMENT FOR OVERDOSE, AND ANTIDOTE(S):

Not Applicable

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS:

Store below 30°C. Keep in airtight container.

PACKING / PACK SIZES:

Box of 100 x 10's and 10 x 10's Blister.

SHELF LIFE:

3 years

Manufacturer and Product Registration Holder:

SM PHARMACEUTICALS SDN BHD (218620-M)

Lot 88, Sungai Petani Industrial Estate

08000 Sungai Petani,

Kedah Darul Aman,

Malaysia.

SM PHARMACEUTICALS SDN. BHD.**MAGNOMINT TABLET**

(MAGNESIUM TRISILICATE, ALUMINIUM HYDROXIDE AND DIMETHICONE TABLET)

DESCRIPTION:

White round flat surface, bevelled edge tablet with break line on one side. Odour: Mint.

COMPOSITION:

Each tablet contains:

Magnesium Trisilicate ... 250 mg

Aluminium Hydroxide... 120 mg

Dimethicone ... 25 mg

Sodium benzoate 0.08% as preservative

ACTIONS AND MODE OR MECHANISMS OF ACTION:

Antacids – React chemically to neutralize or buffer existing quantities of stomach acid, but have no direct effect on its output. This action results in increased pH value of stomach contents, thus providing relief of hyperacidity symptoms.

Dimethicone – acts in vitro to lower surface tension of gas bubbles. Its relevance to action in vivo is not clearly established.

PHARMACOLOGY (SUMMARY OF PHARMACODYNAMICS AND PHARMACOKINETICS):**Absorption:**

Aluminium – A small amount of the aluminium in aluminium hydroxide is absorbed from the intestine.

Magnesium – Approximately 10% of the magnesium in magnesium hydroxide is absorbed from the intestine.

Excretion:

Antacids – Renal and faecal, 15% to 30% of the salts formed are absorbed and excreted by the kidneys.

Dimethicone – Excreted in the faeces as unchanged drug.

19.07.2022

INDICATIONS:

Hyperacidity-Antacid and dimethicone combinations are indicated for the symptomatic relief of heartburn, sour stomach and/or acid indigestion.

CONTRAINDICATIONS:

Renal function impairment.

SIDE EFFECTS / ADVERSE REACTIONS:

Constipation, mild
Diarrhoea or laxative effect
Nausea or vomiting
Speckling or whitish discoloration of stools (concentrations of fatty acids- salts of aluminium)
Stomach cramps.

PRECAUTIONS / WARNINGS:

Not taking this medication:
-If symptoms of appendicitis or inflamed bowel are present, checking with physician for proper diagnosis.
-Within 1 to 2 hours of other oral medication.

Paediatrics

Antacids should not be given to young children (up to 6 years of age) unless prescribed by a physician.

Since children are not usually able to describe their symptoms precisely, proper diagnosis should precede the use of an antacid. Thus will avoid the complication of an existing condition (e.g., appendicitis) or the appearance of more severe adverse effects.

In very young children who are receiving magnesium-containing antacids, there is risk of hypermagnesemia, especially in dehydrated children in those with renal failure.

Geriatrics

Metabolic bone disease commonly seen in elderly may be aggravated by the phosphorus depletion, hypercalciuria, and inhibition of absorption of intestinal fluoride caused by the chronic use of aluminium-containing antacids.

Pregnancy / Reproduction

First trimester – Antacids have been included in large prospective drug surveillance studies. However, one retrospective case-control study has suggested an association of antacids during the first 56 days of gestation and the occurrence of major and minor congenital abnormalities. There was no individual association of any particular antacid with the increased congenital abnormalities.