

ELAN - FORTE[®] CREAM

Ingredient(s):

Each gram contains:

Tolnaftate	10mg
Chlorhexidine HCl	2mg

Preservative(s):

Methyl Paraben	2mg
Propyl Paraben	1mg

Pharmacodynamics:

1. Elan-Forte Cream contains Tolnaftate as its main ingredient. It is a fungicidal and antibacterial agent.
2. Elan-Forte Cream has antifungal activity against thread fungi such as Trichophyton, Microsporum and Epidermophyton.
3. Elan-Forte Cream may be useful in the control of recent, mild fungal infection of the scalp (Tinea capitis).
4. Elan-Forte Cream does not irritate the skin and mucous membrane. It does not stain and has low toxicity.

Pharmacokinetics:

None.

Indication(s):

Ringworm infection of the skin such as Tinea pedis (athlete's foot), Tinea corporis (ringworm of the body), Tinea cruris (ringworm of the groin) and Tinea capitis (ringworm of the scalp); and Tinea versicolor (white spot).

Dosage and Administration:

Apply topically to the site of the lesion twice daily for 2 to 3 weeks. To enhance the effect it should be applied after washing the affected area with luke-warm water.

Route of Administration:

Topical.

Warnings and Precautions:

1. Avoid contact with eyes.
2. If the patient shows no improvement after 4 weeks of treatment, the diagnosis should be re-evaluated; and if the patient's condition becomes more severe the drug should be withdrawn.
3. Not recommended in children up to 2 years of age except under a doctor supervision.
4. This product contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalized allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. This product should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.
5. If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using this product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interaction with Other Medicaments:

None.

Side Effect(s) / Adverse Reaction(s):

Skin reactions including irritation and pruritus may occur.

Immune system disorders

Frequency not known: Hypersensitivity including anaphylactic shock.

Contraindication(s):

None.

Symptoms and Treatment for Overdosage, and Antidote(s):

No significantly toxic reaction in man by topical application.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description:

A white cream.

Dosage Forms and Packing Available:

Cream in:

Aluminium tube of 5g, 10g, 15g and 20g.

Plastic Jar of 50g

Plastic Jar of 500g and 1kg (for export only)

The information contained in this leaflet is limited. Further information can be obtained from other sources including your doctor or pharmacist.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679
ELO - 10

Date of revision: 21 July 2017