

Maalox® Plus Tablet



Composition:

To provide symptomatic relief of hyperacidity plus alleviation of gas symptoms, each tablet contains:

	Maalox® Plus
Active Ingredients	Per Tablet
Magnesium Hydroxide	200mg
Aluminium Hydroxide	200mg
Simethicone (equivalent to Dimethicone)	26.25mg (25mg)

Pharmacological properties:

Pharmacodynamic properties

Pharmacotherapeutic group: Antacids with antiflatulents; ATC code: A02AF02.

Mechanism of action

The antacid effect takes place in the stomach where, at the appropriate dose, magnesium hydroxide and aluminum hydroxide neutralize gastric acid production. The constipating effect of aluminum hydroxide is counteracted by the laxative effect of magnesium hydroxide. Simeticone has an antiflatulent effect.

Pharmacokinetic properties

Absorption

When administered orally, aluminum hydroxide and magnesium hydroxide react with the hydrochloric acid produced in the stomach to form salts, which are only partly absorbed.

Excretion

Absorbed salts are excreted mainly in the urine.

Indications: As an antacid for symptomatic relief of hyperacidity associated with the diagnosis of peptic ulcer, gastritis, peptic esophagitis, gastric hyperacidity, heartburn, or hiatal hernia. As an antiflatulent to alleviate the symptoms of gas, including postoperative gas pain.

Directions for Use: Two to four tablets, well chewed, four times a day, taken twenty minutes to one hour after meals and at bedtime, or as directed by a physician. Maalox® Plus are not recommended for children.

Adverse Effects: The frequency of the undesirable effects listed below is defined using the following conventions: Common (≥1/100 to <1/10); uncommon (≥1/1 000 to <1/100); rare (≥1/10 000 to <1/1 000); very rare (<1/10 000); not known (cannot be estimated from the available data).

Immune system disorders:

Frequency not known: angioedema, anaphylactic reactions, hypersensitivity reactions, urticaria, pruritus.

Gastrointestinal disorders: Uncommon: diarrhea or constipation. Frequency not known: abdominal pain.

Metabolism and nutrition disorders:

Very rare: hypermagnesemia, which has been observed with prolonged use in patients with kidney impairment. Frequency not known: hyperaluminumemia, hypophosphatemia, in prolonged use or at high doses, or even normal doses of the medicinal product in patients on low-phosphorus diets, or in infants (0 to 24 months), which may result in increased bone resorption, hypercalciuria or osteomalacia.

Patient Warnings: Do not take more than 16 tablets in a 24-hour period or use the maximum dosage for more than two weeks or use if you have kidney disease except under the advice and supervision of a physician. Aluminum hydroxide may cause constipation and magnesium salts overdose may cause hypomotility of the intestine; large doses of this medicinal product may trigger or aggravate intestinal obstruction and ileus in patients with higher risk, such as those with renal impairment, primary constipation, impaired intestinal motility, infants (0 to 24 months) and the elderly. Aluminum hydroxide is not well absorbed from the gastrointestinal tract, and systemic effects are therefore rare in patients with normal renal function. However, excessive doses or long-term use, or even normal doses in patients with low-phosphorus diets or in infants (0 to 24 months), may lead to phosphate depletion (due to aluminum-phosphate binding) accompanied by increased bone resorption and hypercalciuria with the risk of osteomalacia. Medical advice is recommended in case of long-term use or in patients at risk of hypophosphatemia. In patients with renal impairment, plasma levels of aluminum and magnesium tend to increase, which causes hyperaluminumemia and hypermagnesemia respectively. In these patients, long-term exposure to high doses of aluminum and magnesium salts may lead to encephalopathy, dementia, microcytic anemia or it may worsen dialysis-induced osteomalacia. Medical monitoring is required in patients with mild to moderate renal failure. Prolonged use of antacids in patients with mild or moderate renal failure should be avoided. Administration of this medicinal product is contraindicated in patients with severe forms of renal failure. Aluminum hydroxide may be unsafe for patients with porphyria who are undergoing hemodialysis

Due to its composition, Maalox® Plus does not generally affect bowel function. However, in some particularly sensitive patients and at high dosages, accelerated intestinal transit may occur.

ARTWORK INFO PANEL

DESCRIPTION: IST MAALOX PLUS 200MG MY I01

PACK TYPE: LEAFLET

NEW CODE: 891393

OLD CODE: 874622

COUNTRY: MALAYSIA

BRAILLE TEXT: N/A

BRAILLE NUMBER: N/A

NOTES:

VERSION: 1

DATE: 13/06/2023

MOCKUP: [NO]

SIZE MM: 167x315 S

DIE CUT: N/A

TECH. DRAW: 62C0018 v 3.0

PACK. LINE: N/A

MINIMUM FONT SIZE: 9,5 PT

PHARMACODE: 62492

PROCESS PLATES

K

NON PRINT

TECHNICAL INFO

SPOT PLATES

| FRONT |

COPYRIGHT // RENIER PHARMA DESIGN // DO NOT TAMPER

BAR PRINTS 100 mm @ 100%

Maalox® Plus contains approximately 15 mg of sorbitol in each tablet. The additive effect of concomitantly administered medicinal products containing fructose (or sorbitol) and dietary intake of fructose (or sorbitol) should be taken into account. The content of sorbitol in medicinal products for oral use may affect the bioavailability of other medicinal products for oral use administered concomitantly.

About 500 mg of glucose in each tablet: this should be taken into account in patients with diabetes mellitus if they are taking more than 10 tablets a day; patients with rare glucose-galactose malabsorption should not take this medicine.

Sucrose; patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicinal product.

Less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

Pediatric populations

Use of magnesium hydroxide in small children, particularly those with kidney impairment or dehydration, may result in hypermagnesemia.

Contraindications:

- Hypersensitivity to the active substances or to any of the excipients.
- Patients with porphyria.
- Severe renal insufficiency (creatinine clearance less than 30 mL/min).
- Generally contraindicated in children.
- Cachexia.

Drug Interaction: Because aluminium and magnesium salts reduce the gastrointestinal absorption of tetracyclines, the use of Maalox® Plus should be avoided during oral tetracycline therapy.

The use of antacids containing aluminium can reduce the absorption of drugs, in particular H2-antagonists, atenolol, cefdinir, cefpodoxime, chloroquine, tetracyclines, dasatinib monohydrate, diflunisal, digoxin, dexamethasone, bisphosphonates, eltrombopag olamine, elvitegravir, ethambutol, fluoroquinolones, glucocorticoids, indomethacin, iron salts, isoniazide, ketoconazole, levothyroxine, lincosamides, metoprolol, nilotinib, phenothiazine neuroleptics, penicillamines, propranolol, raltegravir potassium, rilpivirine, riociguat, rosuvastatin, sodium fluoride, and antiviral treatments in combination with tenofovir alafenamide fumarate/emtricitabine/sodium bictegravir.

- Polystyrene sulfonate (Kayexalate)
Caution is advised when this medicinal product is used concomitantly with polystyrene sulfonate (Kayexalate) due to the potential risk of reduced effectiveness of the resin in binding potassium, of metabolic alkalosis in patients with renal impairment (reported with aluminum hydroxide and magnesium hydroxide) and of intestinal obstruction (reported with aluminum hydroxide).
 - Aluminum hydroxide and citrates may cause hyperaluminumemia, especially in patients with renal impairment.
- As magnesium hydroxide administration results in urine alkalization, increased excretion of salicylates has been observed in the event of concomitant use.

As a precaution, allow at least 2 hours (4 for fluoroquinolones) to elapse between taking any oral drug and Maalox® Plus. Concomitant use of aluminium hydroxide and citrates may result in increased aluminum levels, especially in patients with renal impairment. Urine alkalization following magnesium hydroxide administration may affect the excretion of some drugs; increased excretion of salicylates has been observed.

Pregnancy and Lactation: There are no available data on Maalox® Plus use in pregnant women. No conclusions can be drawn regarding whether or not Maalox® Plus is safe for use during pregnancy.

Pregnancy

The medicinal product should only be used if absolutely necessary, under direct medical supervision, after having evaluated the expected benefits for the mother compared with the possible risk to the fetus or infant.

Breast-feeding

Because of the limited maternal absorption when taken in accordance with the indicated dosage regimen, aluminum hydroxide and its magnesium salt combinations are considered compatible with breast-feeding.

Effects on ability to drive and use machines:

Maalox® Plus has no influence on the ability to drive and use machines.

Overdose:

There is very limited experience with deliberate overdose. Cases of overdose with aluminum salts are more likely to occur in patients with severe chronic renal impairment, with the following symptoms: encephalopathy, seizures and dementia, hypermagnesemia.

The most frequently reported symptoms of acute overdose with aluminum hydroxide and in combination with magnesium salts include diarrhea, abdominal pain and vomiting.

Large doses of this medicinal product may cause or aggravate intestinal obstruction and ileus in patients at risk. Treatment should be symptomatic and include general supportive measures.

Aluminum and magnesium are eliminated through the urinary route; treatment of magnesium overdose consists of rehydration and forced diuresis. In case of renal function deficiency, hemodialysis or peritoneal dialysis is necessary.

Description: white/pale yellow flat face tablet, engraved with “Maalox” on one side.

Supplied: Maalox® Plus Tablets are available in blister (PVC/Alu) packages of 10 tablets, 40 tablets and 100 tablets. Not all presentations are marketed.

Storage condition: Store below 25°C

Shelf-life: Do not utilize after the expiry date indicated on the packaging.

Keep this and all drugs out of the reach of children.

* Each Maalox® Plus Tablet contains approximately 0.03 mEq sodium per tablet.

Manufactured by:

Sanofi S.r.l.
SS 17 KM 22, 67019 SCOPPITO, ITALY.

Revision Date: March 2023
(based on IT SmPC June 2022)



891393

PH 62492

ARTWORK INFO PANEL				
DESCRIPTION: IST MAALOX PLUS 200MG MY I01		VERSION: 1	PROCESS PLATES	
PACK TYPE: LEAFLET NEW CODE: 891393 OLD CODE: 874622 COUNTRY: MALAYSIA BRAILLE TEXT: N/A BRAILLE NUMBER: N/A		DATE: 13/06/2023 MOCKUP: [NO] SIZE MM: 167x315 S DIE CUT: N/A TECH. DRAW: 62C0018 v 3.0 PACK. LINE: N/A MINIMUM FONT SIZE: 9,5 PT PHARMACODE: 62492		
			SPOT PLATES	
NOTES:				
COPYRIGHT // MENIER PHARMA DESIGN // DO NOT TAMPER		BAR PRINTS 100 mm @ 100%		
BACK				