

Item Code	: 204300 - 08 draft 09 Jun 2025
PM Code	: INS.P - AOS
Measurement(mm) ± 2mm	: 180 (L) x 150 (W)
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

AXOL LIQUID

Ambroxol, a metabolite of Bromhexine, is a mucolytic agent. It is capable of increasing sputum volume and reducing its viscosity, improving breathing and reducing cough with no influence on gastric acid secretion.

Ingredient(s):

Each ml contains:

Ambroxol Hydrochloride 3mg
Ethanol (as 96%) 2% v/v

Preservatives:

Methyl Paraben 0.06 % w/v
Propyl Paraben 0.03 % w/v

Pharmacodynamics:

Ambroxol is a mucus-modifying drug; it enhances both bronchial glandular cell secretions and the production of alveolar type II pneumocyte surfactant. It is also stimulates ciliary motility and accelerates mucociliary transport.

1. Axol is a " mucus-modifier ", the active compound of which is Ambroxol. Several human studies showed that Ambroxol affects the secretomotor activity of serous glands, facilitates the repair of bronchial epithelium and accelerates mucociliary transport and clearance.
2. Ambroxol, a metabolite of bromhexine has been found to stimulate the synthesis of lung surfactant, an agent that may play a role in airway mucus transport acting as a lubricant.
3. In several studies, Ambroxol also produced statistically significant symptomatic improvement, measured as difficulty in expectoration, coughing, presence of dyspnea and the auscultatory sign as compared to controls.

Pharmacokinetics:

Absorption after oral administration of Ambroxol was found to be fast and complete. Elimination half-life in the blood was estimated as 20-25 hrs in man. In man and rabbit, the radio-active labeled substance was excreted almost completely into the urine.

Indication(s):

Mucolytic agent used in the treatment of respiratory disorders associated with viscid mucus.

Dosage and Administration:

Adults and children over 10 years old : 10ml 3 times daily.
Children, 5-10 years old : 5ml 2 to 3 times daily.
2-5 years old : 2.5ml thrice daily.

Under 2 years old : 2.5ml twice daily.
Childrens' dose may also be based on body weight : 1.2-1.6mg/kg/day.

Route of administration:

Oral administration

Contraindication(s):

It is contraindicated in patients who have shown hypersensitivity to this drug.

Side Effect(s) / Adverse Reaction(s):

Gastric discomfort, gastralgia, diarrhea, nausea, vomiting, constipation and dyspepsia. Other side effects such as skin rash, pruritus and erythema have also been reported.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Pregnancy and Lactation:

The benefits of the drug in pregnant women during first trimester should be weighed against a possible risk to the fetus.

Precaution(s) / Warning(s):

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Symptoms and Treatment for Overdosage and Antidote(s):

There is no specific antidote for its overdose; therefore, management of the patient should consist of symptomatic and supportive therapy.

L
180mm
(±2mm)

W
150mm
(±2mm)

Item Code	: 204300 - 08 draft 09 Jun 2025
PM Code	: INS.P - AOS
Measurement(mm) ± 2mm	: 180 (L) x 150 (W)
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

Drug Interaction(s):

Not applicable.

Product Description:

Clear, colorless solution to yellowish solution.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Packing(s):Plastic bottle of 60ml, 100ml and 120ml.
Plastic bottle of 1L and 4L (for export only).

暢痰液

成份:

每毫升含:

Ambroxol Hydrochloride 3mg

Ethanol (as 96%) 2% v/v

Preservatives:

Methyl Paraben 0.06 % w/v

Propyl Paraben 0.03 % w/v

適應症:

化痰。

暢痰液能調節粘膜分泌物，增進粘液纖毛運送機轉，促使排除呼吸道內粘稠且阻塞的痰液。同時能抑制咳嗽反射作用，舒緩疲憊刺激性的咳嗽。

用法、用量:

成人及10歲以上兒童 : 每次10ml, 每天3次。

5-10 歲兒童 : 每次 5ml, 每天2次。

2-5 歲兒童 : 每次2.5ml, 每天3次。

2 歲以下兒童 : 每次2.5ml, 每天2次。

兒童劑量亦可按每公斤體重每天1.2-1.6mg。

AXOL LIQUID

Bahan Aktif:

Setiap ml mengandungi:

Ambroxol Hydrochloride 3mg

Ethanol (as 96%) 2% v/v

Pengawet:

Methyl Paraben 0.06 % w/v

Propyl Paraben 0.03 % w/v

Indikasi:Digunakan untuk merawat gangguan pernafasan
berhubung dengan mukus tebal.**Dos:**Dewasa dan kanak-kanak lebih 10 tahun: 10ml 3 kali
sehari.

Kanak-kanak 5 -10 tahun : 5 ml 2 - 3 kali sehari.

2 - 5 tahun : 2.5 ml 3 kali sehari.

bawah 2 tahun : 2.5 ml 2 kali sehari.

Dos untuk kanak-kanak juga boleh diberi mengikut berat
badan : 1.2 - 1.6 mg / kg sehari.

Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 19 Mar 2018

W
150mm (±2mm)

L
180mm
(±2mm)