

DYNAVATE CREAM

MAL19950042AZ

DESCRIPTION:

CREAM

Colour : White
Fragrance : Baby Jo

CONTENT:

Betamethasone 17-Valerate equivalent to Betamethasone 0.1% w/w
(Fragrance: Baby Jo, Preservative: Chlorocresol 0.1%)

PHARMACODYNAMICS:

Corticosteroids diffuse across cell membranes and complex with specific cytoplasmic receptors. These complexes then enter the cell nucleus, bind to DNA and stimulate transcription of messenger RNA (mRNA) and subsequent protein synthesis of various enzymes thought to be ultimately responsible for anti-inflammatory effects to topical application of corticosteroid.

PHARMACOKINETICS:

Corticosteroids when administered by topical application, particularly under an occlusive dressing or when the skin is broken, sufficient corticosteroid may be absorbed to give systemic effects. Corticosteroids in the circulation are extensively bound to plasma proteins, mainly to globulin and less so to albumin. The corticosteroid-binding globulin has high affinity but low binding capacity, which the albumin has low affinity but large binding capacity. Only unbound corticosteroid has pharmacological effects or is metabolised. The synthetic corticosteroids are less extensively protein bound than hydrocortisone (cortisol). They also tend to have longer half-lives. Corticosteroids are metabolised mainly in the liver but also in the kidney, and are excreted in the urine. The slower metabolism of the synthetic corticosteroids with their lower protein-binding affinity may account for their increased potency compared with the natural corticosteroids.

INDICATION:

For the relief of inflammatory manifestations of corticosteroid-responsive dermatoses.

RECOMMENDED DOSE:

Apply thinly to the affected parts two to three times daily.

CONTRAINDICATION:

Rosacea, acne vulgaris, perioral dermatitis, viral infections, bacteria and fungal infections and hypersensitivity to the preparation. It is also contraindicated in patients with diabetes mellitus, tuberculosis and glaucoma.

WARNING AND PRECAUTIONS:

Topical applications of corticosteroid should not be made with an occlusive dressing to large areas of the body because of the increased risk of systemic toxicity and should not, in general, be used in the presence of infection. Occasionally they may be used with the addition of a suitable antimicrobial substance in the treatment of infected skin but there is a risk of sensitivity reactions occurring. Corticosteroid should not be applied to ulcers of the leg and long-term topical use is best avoided, especially in children.

PREGNANCY AND LACTATION:

There is inadequate evidence of safety in human pregnancy. Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development including cleft palate and intra-uterine growth retardation. There may therefore be a very small risk of such effects in the human foetus.

SIDE EFFECTS/ADVERSE REACTIONS:

The following local adverse reactions are reported infrequently with topical corticosteroids but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence; burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae and miliaria.

SYMPTOMS AND TREATMENT OF OVERDOSE:

For topical application of corticosteroids, problems of overdosage are unlikely to be encountered. For chronic topical overdose, treatment is symptomatic, supportive, and consists of discontinuance of topical adrenocorticoid therapy. Gradual withdrawal of the preparation may be necessary. Consult the doctor immediately if any unusual serious reactions happened.

PACKING/PACK SIZE(S):

Aluminium tubes of 15 g and 50 x 15 g.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
FOR EXTERNAL USE ONLY**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA

PI1PE D002-03
Latest Revision: June 2017