

SALCODYL SYRUP

MAL19950247AZ

DESCRIPTION:

SYRUP

Colour : Red

Flavour : Rose

CONTENT:

Each 5 ml contains:

Salbutamol Sulphate equivalent to Salbutamol 1 mg

Guaiphenesin..... 50 mg

Preservatives : Methyl Paraben 0.15% w/v &

Propyl Paraben 0.02% w/v.

Colouring Agent : Raspberry Red

Flavouring Agent : Rose

PHARMACODYNAMICS:

Salbutamol is a directly acting sympathomimetic amine. Its main action is on the adrenergic receptors in the bronchi and the respiratory tract rather than on the cardiac receptor. For this reason it induces bronchodilation and inhibits bronchospasm in doses which do not produce marked cardiac acceleration.

Guaiphenesin has an expectorant action which increases the output of respiratory tract fluid by reducing adhesiveness and surface tension. The increase flow of less viscid secretions promotes ciliary action and facilitates the removal of mucus. This changes a dry unproductive cough to a cough that is more productive and less frequent.

PHARMACOKINETICS:

Salbutamol is rapidly absorbed after oral administration. After an oral dose of 4 to 8 mg, peak plasma concentrations of about 23 ng/ml for unchanged drug and 50 to 100 ng/ml for unchanged drug plus metabolite are attained in 2.5 to 3 hour. Plasma half-life is 2 to 7 hour. Salbutamol is metabolised by first pass metabolism, the reactions and metabolites involved are not yet identified. About 75 to 95% of an oral dose is excreted in the urine and about 4% in the faeces in 3 days. Following oral administration, 50 to 60% of the urinary excreted material is metabolised.

Guaiphenesin is readily absorbed from gastrointestinal tract. It is rapidly metabolised and excreted in the urine.

INDICATION:

Relief of productive and wheezing cough associated with pneumonia, influenza and acute respiratory infections.

RECOMMENDED DOSE:

Adults : 2 to 4 teaspoonfuls (10 ml - 20 ml) 2 to 3 times daily.

Children 2-6 years : 1 to 2 teaspoonfuls (5 ml - 10 ml) 2 to 3 times daily.

6-12 years : 2 teaspoonfuls (10 ml) 2 to 3 times daily.

CONTRAINDICATION:

Salbutamol is contraindicated in patients with hypertension, myocardial insufficiency and hyperthyroidism.

WARNING AND PRECAUTIONS:

Salbutamol should be used with caution in patients suffering from thyrotoxicosis and in patients with diabetes mellitus, serious cardiovascular disorders or hypertension. Safety & effectiveness have not been established in children below 2 years of age.

Tocolysis: Serious adverse reactions including death have been reported after administration of Salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increase fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

DRUG INTERACTION:

Salbutamol should be administered with extreme caution in patients being treated with MAOI or Tricyclic Antidepressant because of the action of Salbutamol on vascular system may be potentiated.

Salbutamol should not be administered with non-selective beta-adrenoreceptor blocking drugs such as propranolol or oxprenolol. There are also risks of hypokalaemia if high doses of Salbutamol given with corticosteroids, diuretics or theophylline.

PREGNANCY AND LACTATION:

Risk and benefit should be considered when used during pregnancy.

SIDE EFFECTS/ADVERSE REACTIONS:

Salbutamol may cause palpitation, tachycardia and muscle tremors. Guaiphenesin may cause gastrointestinal discomfort and drowsiness. Very high doses may cause nausea and vomiting. Potentially serious hypokalemia may result from β_2 -agonist therapy. In young children 2-6 years of age, some adverse reactions were noted more frequently than in adults and older children.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Salbutamol: Symptoms of overdosage include muscle tremor, flushing, agitation, palpitations, sinus tachycardia and hypokalaemia. Treatment includes gastric lavage or emesis and the use of beta-adrenoreceptor blocking agents, usually propranolol.

Guaiphenesin: When given by mouth in large doses, Guaiphenesin has a relaxant effect on skeletal muscle similar to that of mephenesin which is closely resembles structurally but this effect is not produced by the doses normally employed in the treatment of cough.

PACKING/PACK SIZE(S):

Plastic bottles of 100 ml and 120 ml.

JAUHI UBAT DARIPADA KANAK-KANAK

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, Malaysia.

PI1PL S001-04

Latest Revision: May 2022