

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION:

Usual adult dose: Topical, two times a day, morning and evening.

Pediatric: Topical, once a day.

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTES:

Acute overdosage with topical application of the cream would not be expected to lead to a life-threatening situation. However, topically applied corticosteroid can be adsorbed in sufficient amounts to produce systemic effects.

PACKING / PACK SIZES:

Aluminium collapsible tubes of 15 grams.

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS:

Store below 30°C. Do not freeze. For external use only. Complete the treatment as prescribed by the physician.

SHELF LIFE: 3 years

NAME AND ADDRESS OF MANUFACTURER AND PRODUCT REGISTRATION HOLDER:

SM PHARMACEUTICALS SDN BHD (218620-M)

Lot 88, Sungai Petani Industrial Estate

08000 Sungai Petani

Kedah Darul Aman, Malaysia.

SM PHARMACEUTICALS SDN. BHD.

FLUROCOMB CREAM

(Clotrimazole and Triamcinolone Acetonide Cream)

DESCRIPTION:

A smooth, soft, homogenous cream, white in colour and has slightly waxy odour.

COMPOSITION:

Clotrimazole 10 mg (1% w/w)

Triamcinolone Acetonide 1 mg (0.1% w/w)

ACTIONS AND MODE OR MECHANISMS OF ACTION:

Clotrimazole

Fungistatic; may be fungicidal, depending on concentration; inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell wall membrane and altering its permeability; as a result, loss of essential intracellular elements may occur; also inhibits biosynthesis of triglycerides and phospholipids by fungi; inhibits oxidative and peroxidative enzyme activity, resulting in intracellular buildup of toxic concentration of hydrogen peroxide which may contribute to deterioration of subcellular organelles and cellular necrosis. In *Candida albicans*, inhibits transformation of blastospores into invasive mycelial form.

15.04.2022

Triamcinolone Acetonide

Triamcinolone diffuse across cell membranes and complex with specific cytoplasmic receptors. These complexes then enter the cell nucleuc, bind to DNA (chromatin), and stimulate transcription of messenger RNA (mRNA) and subsequent protein synthesis of various enzymes thought to be ultimately responsible for anti-inflammatory effects of topical application of adrenocorticoids.

PHARMACOLOGY (SUMMARY OF PHARMACODYNAMICS AND PHARMACOKINETICS):***Clotrimazole***

Absorption: Dermal penetration; minimal systemic absorption.

Triamcinolone Acetonide

Absorbed systemically; occlusion, use over extensive areas, or prolonged used increases absorption of topical dosage forms.

INDICATIONS:

Inflammatory fungal skin infections and as a secondary agent in topical treatment of cutaneous candidiasis.

CONTRAINDICATIONS:

Patients hypersensitive to active ingredients or any other ingredients of the cream.

Tubercular of the skin including vaccini, varicella and herpes simplex.

SIDE EFFECTS / ADVERSE REACTIONS:

Skin rash, pain, redness, or pus-containing blisters. Burning, itching, blistering or peeling not present before therapy.

PRECAUTIONS / WARNINGS:

Don't use tight-fitting diapers or plastic pants on child if the diaper area is being treated with this medication. Improvement of condition, with relief of pruritus, usually occurs within first week of therapy. When this medication is used in the treatment of candidiasis, occlusive dressings should be avoided since they provide conditions which favour growth of yeast and release of its irritating endotoxin. Systemic absorption of topical corticosteroids has produced reversible hypothalamic, pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia and glucosuria in some patients.

Pregnancy

Topical adrenocorticoid - containing preparations should not be used on extensive surface areas, in large amounts or for prolonged periods of time in pregnant patients. Problems in humans have not been documented.

Lactation

Safety of the use of topical adrenocorticoids during lactation has not been established.

Drug interactions

There are no known drug interactions or physical incompatibilities between Flurocomb Cream and other drugs.