

pharmaniaga®
CHLORHEXIDA 5 SOLUTION
Chlorhexidine Gluconate 5% w/v

DESCRIPTION

A clear reddish solution without any foreign particles and with lavender odour.

COMPOSITION

Each 100mL of the solution contains 5g of Chlorhexidine gluconate.

PHARMACODYNAMICS

Chlorhexidine is a bisbiguanide antiseptic that is bactericidal (kills bacteria) or bacteriostatic (inhibits bacterial growth) against a wide range of Gram-positive and Gram-negative bacteria.

Chlorhexidine kills bacteria by binding cationic molecule to negatively charged bacterial cell walls and microbial complexes leading to precipitation of cytoplasmic contents of bacterial cell wall which results in cell death. At lower concentration, it acts as a bacteriostatic by alteration of bacterial cell osmotic equilibrium and leakage of potassium and phosphorus.

PHARMACOKINETICS

Chlorhexidine is cationic in nature as it binds strongly to skin, mucosa and other tissues and thus very poorly absorbed.

INDICATIONS

Chlorhexida 5 Solution is used for general antiseptic purposes such as for pre-operative skin disinfection and hand washing, disinfection of wounds, burns or other skin damage or disorders, emergency instrument disinfection and storage of clean instruments.

RECOMMENDED DOSE

Solution for the required strengths should be freshly diluted and avoid contamination.

For the pre-operative skin disinfection and handwashing:

Use a concentration of 0.5% in alcohol (70%).

For disinfection of wounds, burns or other skin damage:

Use a concentration of 0.05% aqueous solution.

For the emergency disinfection of clean instruments:

Immerse for 2 minutes in 0.5% solution in alcohol (70%)

For storage and disinfection of clean instruments:

Immerse for 30 minutes in 0.05% aqueous solution

ROUTE OF ADMINISTRATION

For topical use only.

CONTRAINDICATIONS

Hypersensitivity including anaphylactic shock.

WARNINGS AND PRECAUTIONS

For external use only

Keep this medicine out of the reach of children.

Chlorhexida 5 Solution contains chlorhexidine.

Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare.

Chlorhexida 5 Solution should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine. If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product.

Appropriate therapeutic countermeasures must be instituted as clinically indicated.

INTERACTION WITH OTHER MEDICAMENTS

Incompatible with soaps or other anionic agents.

PREGNANCY AND LACTATION

No effects during pregnancy and lactation are anticipated, since systemic exposure to chlorhexidine gluconate is negligible.

ADVERSE EFFECTS / UNDESIRABLE EFFECTS

Skin and subcutaneous tissue disorder. Dermatitis, pruritus, erythema, eczema, rash, skin irritation, blisters
Immune system disorder

Frequency not known: Hypersensitivity including anaphylactic shock

SYMPTOMS AND TREATMENT OF OVERDOSE

Chlorhexidine gluconate taken orally is poorly absorbed into the blood. Accidental ingestion may require gastric lavage using milk, egg white, gelatine or mild soap. Employ supportive measures as appropriate. Haemolysis has been reported following ingestion or accidental intravenous administration. Haemolysis may necessitate blood transfusion.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

No effects on ability to drive and machine use.

STORAGE CONDITIONS

Store below 30°C, away from direct light. Keep container tightly closed. Susceptible to contamination; take precaution to prevent contamination.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PACK SIZE

Plastic Containers of 500mL & 1Litre.

KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 05

Revision Date : 06 Nov 2024

IL-176.05 PRODUCT REGISTRATION HOLDER AND MANUFACTURER

IDAMAN PHARMA MANUFACTURING SDN BHD (200401023395)

LOT 24 & 25, JALAN PERUSAHAAN LAPAN,

BAKAR ARANG INDUSTRIAL ESTATE, 08000 SUNGAI PETANI,

KEDAH DARUL AMAN, MALAYSIA.