

REXOM™

BROMHEXINE ELIXIR 4MG/5ML

Name and Strength of Active Ingredient(s)

Each 5ml contains :

Bromhexine HCl BP 4 mg

Preservative :

Sodium Benzoate 0.2% w/v

Product Description

Colourless liquid with strawberry flavour.

Pharmacodynamics

Bromhexine is an expectorant/mucolytic agent. Following oral administration, Bromhexine has increased sputum volume and reduced the viscosity of bronchial secretions in chronic bronchitis patients.

Pharmacokinetics

Bromhexine HCl is rapidly absorbed from the gastrointestinal tract and undergoes extensive first-pass metabolism in the liver. It is widely distributed to body tissue. About 85 to 90% of a dose is excreted in the urine mainly as metabolites. Ambroxol is a metabolite of bromhexine. Peak plasma concentrations occur after about 1 hour following oral administration.

Indications

Secretolytic therapy in acute and chronic bronchopulmonary diseases associated with abnormal mucus secretion and impaired mucus transport.

Recommended Dose

Under 2 years: 1.25ml, 3 times a day.

2 to 6 years: 2.5ml, 3 times a day.

6 to 12 years: 5ml, 3 times a day.

Route of Administration

Oral

Contraindications

Bromhexine Elixir 4mg/5ml should not be used in patients known to be hypersensitive to Bromhexine or other components of the formulation.

Warnings and Precautions

Patients being treated with Bromhexine should be notified of an expected increase in the flow of secretions. Bromhexine should be used with care in patients with a history of peptic ulcer disease. Care is also advisable in asthmatic patients.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Interactions With Other Medicaments

Bromhexine may increase the concentration of concurrently administered antibiotics in bronchial secretions. No clinically relevant interactions with other medications have been reported.

Pregnancy and Lactation

Pregnancy:

There is no data on the use of Bromhexine in pregnant women. The effects, if any, on the developing fetus are unknown. Until more information is available, Bromhexine should only be used during pregnancy if the maternal condition justifies the potential risk to the fetus.

Lactation:

Available evidence and/or expert consensus is inconclusive or is inadequate for determining infant risk when used during breastfeeding. Weigh the potential benefits of drug treatment against potential risks before prescribing this drug during breastfeeding.

Side Effects

Gastrointestinal side effects may occur occasionally with bromhexine and a transient rise in serum aminotransferase values has been reported. Other reported adverse effects include headache, dizziness, sweating and skin rashes.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Overdose and Treatment

No symptoms of overdose have been reported in man to date. If they occur, supportive and symptomatic treatment should be provided.

Storage Conditions

Keep container tightly closed. Store below 30°C. Protect from heat and light.

Keep out of reach of children. Jauhi dari kanak-kanak.

Dosage Forms

Oral liquid

Pack Size

60ml, 90ml, 100ml and 120ml bottle.

3600ml bottle (for export only).

Registration Number

MAL19962479XZ

Manufactured by & Product Registration Holder

KCK PHARMACEUTICAL INDUSTREIS SDN. BHD.

Plot 61, Bayan Lepas Industrial Park Phase 4, Lintang Bayan Lepas 1,
11900 Bayan Lepas, Pulau Pinang, Malaysia.

Date of Revision of Package Insert

3 July 2021

PI0012-3