
MIST. CARMINATIVE

Each 5ml contains:

Light Magnesium Carbonate 150mg
Sodium Bicarbonate250mg
Ethanol content.....2.6% v/v

Preservatives

Benzoic Acid.....0.10% w/v

Description:

Product Description

White to off-white suspension with tincture cardamom aromatic flavour.

Pharmacodynamics

Magnesium carbonate serves as an antacid to neutralise stomach acidity by forming soluble magnesium chloride and carbon dioxide with longer lasting effect due to its slow onset of action.

Sodium bicarbonate is a short acting antacid which neutralizes the acidity by producing sodium chloride and carbon dioxide to provide a sense of relief.

Pharmacokinetics

Neutralisation of acidity in stomach by magnesium carbonate produces soluble magnesium chloride and carbon dioxide in the stomach whereby magnesium is absorbed and excreted rapidly via renal in the urine.

Sodium bicarbonate neutralises gastric acid with the production of carbon dioxide whereby bicarbonate ions that are not involved in the reaction is absorbed. In the absence of bicarbonate insufficiency in plasma, the ions are excreted in the urine which is depicted as alkaline followed by diuresis.

Indications

Heartburn, for rapid relief of dyspepsia.

Recommended Dose

Adult: 10-20ml to be taken three times a day

Route of Administration

Oral.

Contraindications

Generally, contraindicated in patients that are hypersensitive to antacids or with hypophosphataemia.

Magnesium carbonate is contraindicated in patients with heart block and severe renal impairment

Sodium bicarbonate is contraindicated in patients with controlled sodium intake, heart failure, hypertension, renal failure, cirrhosis, diarrhoea.

Warning and Precautions

Magnesium carbonate should be cautiously given to patients with less severe degree of cardiac or renal impairment and should be monitored for clinical signs of excess magnesium.

Sodium bicarbonate should be given extremely cautiously to patients with metabolic or respiratory alkalosis, hypocalcemia, hypochlorhydria, heart failure, oedema, renal impairment, hypertension, eclampsia or aldosteronism.

Interactions with Other Medicaments

Oral magnesium salts can possibly decrease the absorption of tetracyclines and bisphosphonates and doses should be separated by a number of hours.

Sodium bicarbonate enhances the lithium excretion and increased renal clearance of acidic drugs such as salicylates, tetracyclines and barbiturates. Sympathomimetics and stimulants prolongs the half life of basic drugs enhancing toxicity.

Pregnancy & Lactation

Magnesium carbonate is not recommended during pregnancy unless fetal heart rate is monitored and usage within 2 hours of delivery should be avoided as magnesium is reported to cross the placenta.

No clinical data on pregnancy nor neonates are reported for sodium bicarbonate. However, it is advisable to cautiously prescribe to pregnant women.

Side Effects

Magnesium carbonate may cause diarrhoea, hypermagnesia, manifesting symptoms like nausea, vomiting, flushing of skin, thirst, hypotension, drowsiness, confusion, muscle weakness.

Sodium bicarbonate can cause stomach cramps, belching and flatulence.

Symptoms and Treatment of Overdose

Symptoms of overdose are high possibilities of hypermagnesia, hypercalcaemia, hypernatremia and hyperphosphatemia

Treatment of overdose to give calcium salts via intravenous administration for hypermagnesia, to give plenty of salt free liquids for the treatment of hyponatremia, and to remove the source of calcium as a measure to overcome hypercalcaemia.

Storage Condition

Store below 30°C. Keep the container tightly closed. Protect from light.

Presentation:

60ml, 90ml, 100ml, 120ml, and 240ml.

Registration Number:

MAL19962484X

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

19 June 2026