
REXOMTM WINA CALAMINE LOTION

Compositions:

Calamine 6.8 % w/v
Zinc Oxide 6.8 % w/v
Ethanol 1.43 % v/v

Product Description:

Off-white to light pink lotion.

Pharmacodynamics:

Calamine has mild astringent and antipruritic actions on the skin and is used to relieve the discomfort of dermatitis as it produces a cooling effect and helps to divert attention away from the itching and thus giving relief from skin irritations.

Zinc oxide is applied externally as a mild astringent for the skin.

Zinc oxide has soothing and protective properties and is used in topical preparations for eczema, slight excoriations, wounds and haemorrhoids. It also reflects ultraviolet radiation and can be used as a physical sunscreen.

Pharmacokinetics:*Absorption*

Calamine and Zinc Oxide are not absorbed through the skin.

Indication:

For the relief of minor skin rashes and irritation.

Recommended Dosage:

Wash and dry affected area before each application.

Apply sparingly to the affected parts and surrounding skin areas twice or three times daily.

Repeated applications may be administered as needed.

****Lotion:** Shake well before use. Lotion should be applied with a cotton ball/pledget moistened with the lotion. Then the medication should be allowed to dry on the skin.

****Extremes of temperature, wearing tight fitting nappies or non-breathable clothing such as polyester or nylon can make itching worse.**

Mode of Administration:

Topical

Contraindications:

Hypersensitivity to any ingredients/components in the formulation.

Warning and Precautions:

For external use. Not to be taken.

Avoid contacts with the eyes and other mucous membranes such as mouth, nose and genital region. Not to be applied to abraded, blistered, inflamed, raw or oozing areas of the skin.

Do not use on open wounds or burns.

Discontinue use if rash, irritation, burning sensation or sensitivity develops/occurs. If symptoms persist, consult your pharmacist or doctor.

Interactions with Other Medicaments:

No clinically significant interactions known.

Pregnancy and Lactation:

The safety of calamine and zinc oxide during pregnancy and lactation has not been established but use during these periods is not considered to constitute a hazard.

Adverse Effects:

Calamine and zinc oxide is well-tolerated except for occasional dryness following and excessive use. Occasional hypersensitivity reactions such as itchiness in sensitive individuals.

Overdose and Treatment:

Overdose arising from the topical application is extremely remote. As the ingredients are inert and very poorly absorbed, effects if any arising from accidental overdose (orally) is unlikely to be life threatening.

If ingested symptoms of overdose include gastritis, nausea, vomiting, epigastric discomfort and diarrhea.

Treatment consists of symptomatic measures only.

Maintain adequate hydration if vomiting or diarrhoea becomes extensive. Milk or antacids may be used to allay gastritis in the event of oral ingestion.

Effects on Ability to Drive and Use Machine:

Unlikely to impair a patient's ability to drive or use machinery.

Storage Conditions:

Keep container tightly closed. Protect from light. Store below 30°C.

Shelf Life:

36 months

Pack Size:

100ml and 120ml bottle.

Registration Number:

MAL19962491XZ

Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Manufactured by:

KCK Pharmaceutical Industries Sdn. Bhd.

No. 5 & 7 Hala Beringin,
Diamond Valley Industrial Park, 11960 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

27 May 2024