

YSP SALBUTAMOL SYRUP 2mg/5ml

Ingredient(s):

Each 5ml contains:

Salbutamol Sulfate 2.41mg
(eq. to Salbutamol base 2mg)

Preservative:

Sodium Benzoate 0.4%W/V

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):**Pharmacodynamic:**

Salbutamol is a potent beta-adrenergic stimulant free from undesirable cardiovascular actions. From a new series of potent beta-adrenergic stimulants, Salbutamol was chosen for development because of its highly selective action on bronchial muscle, its lack of effect on heart muscle and its long duration of action.

Pharmacokinetic:

Salbutamol administered intravenously has a half-life of 4 to 6 hours and is cleared partly renally and partly by metabolism to the inactive 4'-O-sulphate (phenolic sulphate) which is also excreted primarily in the urine. The faeces are a minor route of excretion. The majority of a dose of salbutamol given intravenously, orally or by inhalation is excreted within 72 hours. Salbutamol is bound to plasma proteins to the extent of 10%.

After oral administration, salbutamol is absorbed from the gastrointestinal tract and undergoes considerable first-pass metabolism to the phenolic sulphate. Both unchanged drug and conjugate are excreted primarily in the urine. The bioavailability of orally administered salbutamol is about 50%.

Indication(s):

Indicated for the relief of bronchospasm in bronchial asthma of all types, chronic bronchitis and emphysema.

Dosage and Administration:

Usual adults dose and children above 14 years of age:

Oral 5ml to 15ml, 3-4 times a day initially, the dosage may increase as needed and tolerated up to a maximum of 20ml 4 times a day.

Usual pediatric dose:-

Children up to 2 years of age:- Dosage has not been established.

Children 2 to 6 years of age:- Oral, 0.1mg/kg body weight 3 times a day initially, the dosage being increase as needed and tolerated up to 0.2mg/kg body weight, not to exceed 10ml 3 times a day.

Children 6 to 14 years of age:- Oral, 5ml 3-4 times a day initially, the dosage may increase as needed and tolerated up to a maximum of 60ml a day in divided doses.

Route of administration: Taken orally

Contraindication(s):

It should not be used for threatened abortion during the first and second trimester of pregnancy.

Salbutamol oral preparations are contraindicated in patients with a history of hypersensitivity to any of their components.

Side Effect(s) / Adverse Reaction(s):

The only reported side effect of significance is a fine tremor of skeletal muscle which occurs in some patients; usually the hands are most obviously affected. This effect is dose related and is caused by a direct action on skeletal muscle: It is common to all beta-adrenergic stimulants. A few patients feel tense, this is also due to the effect on skeletal muscle and not to direct CNS stimulation. Nervousness, irregular heartbeat, drowsiness, muscle cramps or twitching or unusual taste may occur.

With dose higher than those recommended or in patients who are unusually sensitive to beta-adrenergic stimulants, peripheral vasodilation and a compensatory small increase in heart rate may occur. Occasionally headaches have been reported.

Hypersensitive reactions including angioedema, urticaria, bronchospasm, hypotension and collapse have been reported very rarely. Potentially serious hypokalaemia may result from beta 2-agonist therapy.

Precaution(s) / Warning(s):

1. Salbutamol Sulfate should be used with caution in patients with hyperthyroidism, diabetes mellitus, serious cardiovascular disorders and hypertension.
2. It should be administered cautiously to patients suffering from thyrotoxicosis.
3. Tocolysis: Serious adverse reactions including death have been reported after administration of salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

Pregnancy and Lactation:

Administration of the drug during pregnancy and lactation should only be considered if the expected benefit to the mother is greater than any possible risk to the fetus/neonate.

Drug Interaction(s):

The concomitant use of Salbutamol and other oral sympathomimetic agent is not recommended since such combined use may lead to deleterious cardiovascular effects. Such concomitant use, however, should be individualized and not given on a routine basis. If regular co-administration is required, then alternative therapy should be considered.

Salbutamol should be administered with extreme caution to patients being treated with monoamine oxidase inhibitors or tricyclic antidepressants because the action of Salbutamol on the vascular system may be potentiated.

Beta-receptor blocking agents and Salbutamol inhibit the effect of each other.

Salbutamol and non-selective beta-adrenergic drugs such as propranolol should not be prescribed together.

Symptoms and Treatment for Overdosage, and Antidote(s):

The expected symptoms with overdosage are those of excessive beta-stimulation and /or occurrence or exaggeration of any of the symptoms e.g. seizures, angina, hypertension or hypotension, tachycardia, arrhythmias, nervousness, headache, tremor, dry mouth, palpitation, nausea, dizziness, fatigue, malaise and insomnia.

Hypokalaemia may also occur.

Treatment consists of discontinuation of Salbutamol together with appropriate symptomatic therapy.

Dialysis is not appropriate treatment for overdosage of Salbutamol syrup.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Description(s):

A clear, colorless to yellowish solution with strawberry flavor.

Dosage Forms and Packaging Available:

Plastic bottle of 60ml, 100ml and 120ml.

Plastic bottle of 1L and 4L(for export only).

YSP SALBUTAMOL SYRUP 2mg/5ml

Bahan Aktif:

Setiap 5ml mengandungi:

Salbutamol Sulfate 2.41mg
(setara dengan Salbutamol base 2mg)

Pengawet:

Sodium Benzoate 0.4%W/V

Indikasi:

Untuk melegakan bronkospasma bagi pelbagai jenis asma bronkial, bronkitis kronik dan emfisema.

Dos:

Dewasa dan kanak-kanak berumur 14 tahun ke atas:-

Dos permulaan 5ml -15ml diberi 3-4 kali sehari dan diubahsuai bila perlu ke maximum 20ml, 4 kali sehari.

Dos kanak-kanak:-

2 tahun ke bawah:- Dos belum diperkenalkan.

2 - 6 tahun:- Dos permulaan 0.1mg/kg berat badan, 3 kali sehari dan diubahsuai bila perlu ke 0.2mg/kg berat badan dan tidak melebihi 10ml 3 kali sehari.

6-14 tahun:- Dos permulaan 5ml, 3-4 kali sehari dan diubahsuai bila perlu ke maximum 60ml sehari diberikan dalam dos terbahagi.



Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679