

Bekunis®

Laxative Dragées

(Enteric coated tablet)

Composition	1 gastric-juice-resistant tablet contains
Active Ingredient	Dry extract of Tinnevelly Senna pods corresponding to 20 mg hydroxyanthracene derivatives calculated as Sennoside B 150 – 220 mg.
Other Ingredients	Polyethylene glycol 6000, Liquid glucose, Microcrystalline cellulose, Polysorbate 80, Acacia gum, Magnesium stearate, Calcium carbonate, Talc, Sucrose, Lactose anhydrous, Titanium dioxide, Stearic acid, Hydrophobic colloidal silica, Methacrylic copolymer, Sodium carboxymethylcellulose, Montanglycolwax.
 Indications	For short term use in cases of constipation of varying causes, conditions in need of easy bowel movements and for clearing or emptying the intestinal tract for diagnostic and therapeutic measures.
Contraindications	Ileus, inflammatory conditions of the intestine (e.g. Crohn's disease, ulcerative colitis) as well as abdominal pains of unknown cause and electrolyte imbalance. Also contraindicated in patients with a hypersensitivity to preparation of Senna. Do not give to children under 12 years of age. Do not use during pregnancy and the lactation period.
Dosage and Mode of Application	If not prescribed otherwise, swallow 1 – 2 tablets with sufficient liquid in the evening. Stimulating laxatives should not be taken over a long period without consulting a physician.
Side Effects	Are dose dependent but rare. a) abdominal pains b) abdominal colic c) tenesmus d) nausea e) belching f) diarrhoea g) may discolour urine or stools. Chronic intake of high doses can lead to hypokalaemia, albuminuria and haematuria.
Precautions	If condition persists for more than one week, please consult a physician. Long term use is not advisable. In chronic cases of constipation, treatment with high fiber diet should be encouraged. Frequent or prolonged use of this preparation may result in dependence towards the product and imbalanced electrolytes. Do not use when abdominal pain, nausea or vomiting are present. Please consult a healthcare practitioner for use beyond 7 days. The shelf life of the product is 3 years from the date of manufacture. Do not use after the expiry date. Store below 25 °C. Jauhkan dari kanak-kanak. Keep out of reach of children.
Product Description	The preparation Bekunis Laxative Dragées is an enteric-coated tablet with the medicinally effective constituent dry extract from Tinnevelly senna pods as well as adjuvants commonly used in the production of sugar-coated tablets.

350331 1406

Pharmacokinetics	The sennosides are prodrugs which are neither split nor absorbed in the upper gastro-intestinal tract. In the colon they are broken down by bacterial enzymes into rhein anthrone. Rhein anthrone is the laxative metabolite. The systemic availability of rhein anthrone is very small. In animal experiments, less than 5 % is discharged in the urine in the form of the oxidised and partly conjugated products rhein and sennidine. The greatest part of the rhein anthrone (> 90 %) is bonded to intestinal content in the faeces and is excreted in the form of polymeric compounds. Small quantities of active metabolites such as rhein find their way into the mother's milk. No laxative effect has been observed in suckling infants. In animal experiments, the presence of rhein in the placenta is extremely low.
Pharmacodynamics	Apart from mucins and flavonoids, senna contains compounds derived from anthracene, the 1,8-dianthrol derivatives. These have a laxative effect. In the case of the sennosides, which are quantitatively predominant, or of their active metabolite in the colon, rhein anthrone, this effect is based mainly on an influencing of colonic motility in the sense of an inhibition of the stationary contractions and a stimulation of the propulsive contractions. This results in an accelerated passage through the intestines and, due to the shortened contact time, in a reduction of the absorption of liquid. In addition, water and electrolytes are discharged through a stimulation of the active secretion of chlorine.
Pregnancy and Lactation	Not to be used in pregnancy and lactation.
Symptoms and Treatment of Overdose	Cases of acute oral laxative intoxication are characterised above all by massive diarrhoea, in some cases accompanied by severe intestinal irritation and in some cases accompanied by massive water and electrolyte losses. This may lead to hypokalaemic and painful muscular spasms. If colics and flatulence are initially prevalent, in some circumstances it may be possible for the clinical picture of acute abdomen to be simulated. Chronic abuse of laxatives is characterised above all by hypokalaemia and often also by inanition. Elementary remedial action: Equalisation of the water-electrolyte balance.
Interactions with other Medicaments	In the case of chronic abuse, a strengthening of the cardiac glycoside effect is possible as a result of hypokalaemia. Simultaneous use with chloramphenicol can significantly reduce the laxative effect of senna.

45 dragees

roha arzneimittel GmbH
Rockwinkeler Heerstr. 100
28355 Bremen, Germany

Date of revision May 2011

Art. 012650 935