

Symptoms and Treatment of Overdose:

The signs and symptoms of overdose are headache, nausea, vomiting, blurred vision, dilated pupils, bot, dry skin, dizziness, dryness of the mouth, difficulty in swallowing, CNS stimulation. Treatment should consist of gastric lavage, emetics and activated charcoal. Barbiturates may be used either orally or intramuscularly for sedation but they should not be used if Acolic syrup with phenobarbital has been ingested. If indicated, parenteral cholinergic agents such as Urecholine (bethanecol chloride) should be used.

Storage Condition:

Keep container tightly closed. Store in a dry place (below 30°C). Protect from light.

Pack Size:

Glass bottle containing 30ml, 60ml, 90ml, 100ml and 120ml.
Glass bottle containing 1000ml for export purpose only.

Product Registration Number:

MAL19972939AZ

Further information can be obtained from your doctor or pharmacist.

ROYCE®

Product Holder / Manufactured by:

ROYCE PHARMA MFG. SDN.BHD. (650435-X)

PT 1663, Nilai Industrial Estate,

71800 Nilai, Negeri Sembilan, Malaysia.

210817

ROYCE®

ACOLIC SYRUP

(Dicyclomine hydrochloride with
Simethicone - Antispasmodic / Antiflatulent)

Presentation:

An orange colour syrup with banana flavour syrup.

Each 5ml contains: Dicyclomine hydrochloride 5mg
Simethicone 50mg

Preservatives:

Methyl paraben 0.1%w/v
Propyl paraben 0.01%w/v
Sodium benzoate 0.1%w/v

Pharmacological Information:

Dicyclomine HCl relieves smooth muscle spasm of the gastro-intestinal tract. Simethicone - an antiflatulence.

Dicyclomine HCl is readily absorbed from the gastro-intestinal tract. It is excreted in the urine, partly as a metabolite.

Indication:

Acolic syrup is indicated for the treatment of functional conditions involving smooth muscle spasm of the gastro-intestinal tract. The commonest of these are: irritable colon (mucous colitis, spastic colon) spastic constipation, infant colic. Acolic syrup can also be used as adjunctive medication in a number of organic gastro-intestinal conditions to relieve associated smooth, muscle, e.g. colitis, diverticulosis and diverticulitis, regional enteritis, gastritis, peptic ulcer, ulcerative colitis, cholelithiasis and cholecystitis (but no biliary colic).

Dosage and Administration:

Not recommended for use in infant under the age of 6 months.

Adults :Two to four 5ml measures 3 or 4 times daily before or after meals.

Children: 2 to 12 years: Two 5ml measures three times daily.

6 months to 2 years: One to two 5ml measures 3 or 4 times daily, 15 minutes before feeds.

Contraindication:

Obstructiveuropathy (for example bladder neck obstruction due to prostatic hypertrophy); obstructive disease of the gastro-intestinal tract (as an achalasia, pyloroduodenal stenosis); paralytic ileus, intestinal atony of the elderly or debilitated patient; unstable cardiovascular status in acute hemorrhage; severe ulcerative colitis; toxic megacolon complicating ulcerative colitis, myasthenia gravis.

Warning and Precautions:

Dicyclomine is not recommended for use in infants under the age of 6 month.

In the presence of a high environment temperature, heat prostration can occur with drug use (fever and heat stroke due to decreased sweating).

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance treatment with this drug would be in appropriate and possibly harmful.

Acolic syrup may produce drowsiness or blurred vision. In this event, the patient should be warned not to engage activities requiring mental alertness such as operating a motor vehicle or other machinery or perform hazardous work while taking this drug.

There are rare reports of infants, 6 weeks of age and under, administered dicyclomine hydrochloride syrup, who have evidenced respiratory symptoms (breathing difficulty, shortness of breath, breathlessness, respiratory collapse, apnea), as well as seizures, syncope, asphyxia, pulse rate fluctuations, muscular hypotonia and coma. The above symptoms have occurred within minutes ingestion and lasted 20 to 30 minutes. The timing and nature of the reactions suggest that they were a consequence of local irritation and/or aspiration rather than a direct pharmacologic effect. Not known deaths or permanent adverse effects have been reported. Acolic syrup should be used with caution in this age group.

Although studies have failed to demonstrate adverse effects of dicyclomine hydrochloride in glaucoma or in patients with prostatic hypertrophy, it should be prescribed with caution in patients known to have or suspected of having glaucoma or prostatic hypertrophy.

Use with caution in patients with:

- Autonomic neuropathy
- Hepatic or renal disease
- Ulcerative colitis. Large doses may suppress intestinal motility to the point of producing a paralytic ileus and the use of this drug may precipitate or aggravate the serious complication of toxic megacolon.
- Hyperthyroidism, coronary heart disease, congestive heart failure, cardiac arrhythmias and hypertension.

Do not rely on the use of the drug in the presence of complication of biliary tract disease. Investigate any tachycardia before giving anticholinergic (atropine-like) drugs since they may increase the heart rate. With overdosage, a curare-like action may occur.

Interactions with Other Medicaments:

Not known drug interaction reported for Dicyclomine hydrochloride.

Pregnancy and Lactation:

Use is not recommended.

Side Effects:

Anticholinergics/antispasmodics produce certain effects which may be physiologic or toxic depending upon the individual patients's response. The physician must delineate these.

Adverse reactions may include xerostomias, urinary hesitancy and retention, blurred vision and tachycardia, palpitations, mydriasis, cycloplegia, increased ocular tension, loss of taste, headache, nervousness, drowsiness, weakness, dizziness, insomnia, nausea, vomiting, impotence, suppression of lactation, constipation, bloated felling, severe allergic reaction or drug idiosyncrasies including anaphylaxis, urticaria and other dermal manifestations, some degree of mental confucion and/or excitement, especially in elderly persons and decreased sweating. With the injectable form there may be a temporary sensation of light headedness and occasionally local irritation.