

**BUVENTOL EASYHALER® 100 mcg/ dose
Inhalation Powder**
**BUVENTOL EASYHALER® 200 mcg/ dose
Inhalation Powder**

Product Description

BUVENTOL EASYHALER is an inhalation powder packed into a multiple dose powder inhaler from which the patient can accurately deliver the individual drug doses. It is for oral inhalation.

BUVENTOL EASYHALER 100 mcg/ dose Inhalation Powder:
The product contains salbutamol sulphate equivalent to 100 mcg of salbutamol per metered dose. The dose, which the patient gets through the mouthpiece of the device is 90 mcg of salbutamol.

BUVENTOL EASYHALER 200 mcg/ dose Inhalation Powder:
The product contains salbutamol sulphate equivalent to 200 mcg of salbutamol per metered dose. The dose, which the patient gets through the mouthpiece of the device is 180 mcg of salbutamol.

The BUVENTOL EASYHALER multiple dose powder inhaler contains 200 doses of finely powdered salbutamol sulphate mixed in a small quantity of lactose. The amount of drug received by the patient (90 or 180 mcg/dose) corresponds to that from salbutamol inhalation aerosol. The dose to be inhaled is released from the container by pressing the device between the thumb and forefinger. This is followed by inhalation through the device. As a result, salbutamol particles are transferred to their target, the lungs. Use of the BUVENTOL EASYHALER does not require coordination of actuation of the dose and inhalation.

Pharmacodynamics

BUVENTOL EASYHALER contains a selective beta₂-adrenergic receptor agonist, salbutamol. Salbutamol acts especially on the bronchi where it relaxes the smooth muscles. The effect of salbutamol is mediated by cyclic AMP. Salbutamol also inhibits the release of histamine and some other mediators in the lungs and accelerates ciliary clearance of the bronchi.

When shifting from freon-based metered dose aerosols to other drug dispensing devices, it is optimal to offer the patient a dispensing device, which does not require adjustment of the dose of the antiasthmatic. It is easiest for the patient to manage a new drug dispenser, which resembles as closely as possible his/her previous aerosol dispenser.

The salbutamol powder dispensed by EASYHALER has been shown to exhibit clinically equivalent bronchodilatation to the optimally used salbutamol aerosol when equal metered doses are administered.

Pharmacokinetics

Inhaled salbutamol acts rapidly, and maximum response is reached in 30 -60 minutes. Following oral inhalation, the fraction that is distributed to the lung (approx. 10 - 25 %) is rapidly seen in the circulation as free unmetabolized drug. The plasma concentrations of inhaled salbutamol are, however, lower than those produced by oral dose. At the doses used in inhalation therapy, the adverse reactions to salbutamol are clearly less than in oral therapy. Salbutamol and its metabolites are rapidly excreted in the urine and faeces with about 80% of the dose being recovered in urine within 24 hours. The elimination half-life of salbutamol is 2.7 to 5.5 hours after oral and inhaled administration.

When shifting from freon-based metered dose aerosols to other drug dispensing devices, it is optimal to offer the patient a dispensing device, which does not require adjustment of the dose of the antiasthmatic. It is easiest for the patient to manage a new drug dispenser, which resembles as closely as possible his/her previous aerosol dispenser.

The salbutamol powder dispensed by EASYHALER has been shown to exhibit clinically equivalent bronchodilatation to the optimally used salbutamol aerosol when equal metered doses are administered.

Indications

Symptomatic treatment of acute asthma attack. Prevention of exercise-induced bronchospasm. Symptomatic treatment of bronchial asthma and other conditions, such as bronchitis and emphysema, associated with reversible airways obstruction.

Contraindications

History of hypersensitivity to any of the components of the product.

Side Effects

Inhaled salbutamol is normally well tolerated. Side effects are rare and they usually disappear with continued treatment. Some patients may experience slight tremor of skeletal muscles and palpitation. These side effects are dose dependent.

Transient nausea, headache, restlessness, and dizziness have been reported. Hypersensitivity reactions including bronchospasms, urticaria, hypotension, and collapse have also been reported. Rare adverse events are dryness and irritation of the mouth, irritation of the throat, muscle cramps, and hypokalemia.

Special Precautions / Warnings

If previously effective dose of salbutamol inhaler fails to give relief for at least three hours, the patient should be advised to contact his doctor for evaluation of the need of additional anti-inflammatory medication. Potentially serious hypokalemia may result from beta₂-agonist therapy.

Particular caution is advised in acute severe asthma as this effect may be potentiated by concomitant treatment with xanthine derivatives, steroids, diuretics and also by hypoxia. Serum potassium levels are recommended to be monitored in such situations.

In patients with diabetes, the blood glucose level should be checked in the beginning of the treatment until the possible slight hyperglycaemic effect of salbutamol is balanced by the treatment of diabetes.

Salbutamol should be administered cautiously to the patients suffering from thyreotoxicosis.

Effects on ability to drive and use machines

BUVENTOL is unlikely to affect ability to drive or to operate machinery.

Drug Interactions

Beta-blockers, such as propranolol, inhibit the effect of salbutamol. Additional adrenergic drugs, monoamine oxidase inhibitors, or tricyclic antidepressants should be used with caution concomitantly with salbutamol because of deleterious cardiovascular effects.

Use in Pregnancy and Lactation

Unnecessary use of the drug should be avoided, particularly during the first trimester of pregnancy. As salbutamol is excreted in breast milk its use in nursing mothers requires careful consideration.

Dosage and Administration

The preparation is intended for oral inhalation only. This preparation is particularly useful for patients who are unable to use metered dose inhaler properly and for patients to whom the use of inhalation aerosol causes irritation of airways. The lowest effective doses of inhaled salbutamol are recommended and instead of regular use it is recommended to use inhaled salbutamol when needed.

Adults: For the relief of acute asthma or before exercise 100-400mcg. The recommended dose for maintenance treatment or prophylactic therapy is 100-400 mcg two to four times a day.

Maximum dose is 2.4 mg/day. Children: In the treatment of episodic asthma or before exercise 100-200 mcg. The recommended dose for maintenance or prophylactic treatment is 100-200 mcg two to four times daily. Maximum dose is 1.2 mg/ day.

Instructions for Use / handling:

Patients have to be instructed to perform a strong and deep inhalation through the EASYHALER device. Patients have to be instructed not to exhale into the device. Illustrated user's instructions for use accompany each package.

Overdosage

Overdose symptoms are those of excessive beta-stimulation, e.g. seizures, angina, hypertension or hypotension, tachycardia with rates up to 200 beats/min, arrhythmias, nervousness, headache, tremor, dry mouth, palpitation, nausea, dizziness, fatigue, and insomnia.

Treatment consists of discontinuation of salbutamol together with appropriate symptomatic therapy. Administer a cardioselective beta-adrenergic blocker (e.g. acebutolol, atenolol, metoprolol), if necessary for cardiac arrhythmias. However, the beta-adrenergic blocker should be used with caution because it could induce severe bronchospasm.

Packaging

BUVENTOL EASYHALER 100 mcg/dose Inhalation Powder: A User Package containing 1 unit of BUVENTOL EASYHALER 100 mcg/dose inhalation powder in the inhaler device, and instruction for use.

BUVENTOL EASYHALER 200 mcg/dose Inhalation Powder: A User Package containing 1 unit of BUVENTOL EASYHALER 200 mcg/dose inhalation powder in the inhaler device, and instruction for use.

Storage: Store in a dry place Store below 30° C. Keep out of reach of children.

Shelf life: 3 years when stored in laminate pouch, 6 months opening of the pouch.

BUVENTOL EASYHALER should not be used after these dates.

Manufactured by
Orion Corporation
Espoo, Finland

Product Registration Holder
ORION PHARMA (MY) SDN. BHD.
E-20-01, Menara SUEZCAP 2,
KL Gateway, No. 2, Jalan Kerinchi, Gerbang Kerinchi Lestari,
59200 Kuala Lumpur, Malaysia

Revision Date:

June 2022

INSTRUCTIONS FOR USE BUVENTOL EASYHALER®

Salbutamol inhalation powder


Read the following instructions before using BUVENTOL EASYHALER inhalation powder. Follow the instructions carefully. With BUVENTOL EASYHALER multidose powder inhaler you can inhale salbutamol to your lungs without needing to inhale any propellants. The powder inhaler contains 200 doses of salbutamol, which is a broncho dilating drug substance used in the prophylaxis and treatment of asthma attacks.

When you first get your EASYHALER

The EASYHALER comes in a foil bag. Do not open the foil bag until you are ready to start using the medicine as it helps to keep the powder dry in the inhaler.

When you are ready to start treatment open the bag and record the date e.g., in your calendar.

Use the inhaler within 6 months of removing from the foil bag.



HOW TO USE CORRECTLY

Step 1: SHAKE

- Remove the dustcap
- Shake the inhaler **3 to 5** times holding it in the **upright position**

SHAKE x 3-5




Important points to remember

- It is important to keep the inhaler in the upright position
- If you accidentally click while you shake the inhaler, empty the powder from the mouthpiece as shown below

Step 2: CLICK

- Keep holding the inhaler upright between your forefinger and thumb
- Press down until you hear a click, and let the inhaler click back again. This releases a dose
- Only click down once**

CLICK x 1




Important points to remember

- The inhaler will not click if the dustcap is still on
- Only click down once
- If you accidentally click more than once, empty the powder from the mouthpiece, see below
- Click to release dose before you inhale, not at the same time
- Keep the inhaler **upright** when you click it and when you inhale the dose. If you tip it, the powder could fall out before you are able to inhale it

Step 3: INHALE

- Keep holding the inhaler upright
- Breathe out normally
- Place the mouthpiece in your mouth between your teeth and close your lips tightly around the mouthpiece
- Take a strong and deep breath in
- Take the inhaler out of your mouth, hold breath **for at least 5 seconds**, then breathe out normally.

INHALE



Important points to remember

- Make sure the whole mouthpiece is well inside your mouth, so that the medication gets into your lungs
- Make sure your lips make a good seal around the mouthpiece
- Do not breathe out into the inhaler. This is important: it could clog up the inhaler. If you breathed out into the inhaler, empty the powder from mouthpiece, see below

If you need to take another inhalation, please repeat the steps 1-3 Shake-Click-Inhale.

After you have used the inhaler:

- Put the dust cap back on the mouthpiece. It stops the inhaler going off by accident.
- After you have taken the dose, rinse your mouth with water, and spit it out and/or brush your teeth.

This may help to stop you getting thrush and becoming hoarse.


How to empty the powder from the mouthpiece

If you click the inhaler by accident, or if you might have clicked it more than once, or if you breathe out into it, empty the mouthpiece.

- Tap the mouthpiece to empty the powder onto a table top, or the palm of your hand.
- Then start again with steps Shake-Click-Inhale.

Cleaning the EASYHALER

Keep your inhaler dry and clean. If necessary you may wipe the mouthpiece of your inhaler with a dry cloth or tissue. Do not use water: the powder in the EASYHALER is sensitive to moisture.



Using the EASYHALER with a protective cover

You may use a protective cover with your inhaler. This helps to improve durability of the product. When you first insert your inhaler in the protective cover make sure the dustcap is on the inhaler as this stops it going off by accident. You can use the inhaler without removing it from the protective cover.

Follow the same instructions as above, **1. Shake – 2. Click – 3. Inhale.**

Remember to:

- Keep the inhaler in the upright position when clicking it
- Replace the dustcap after taking the dose as this stops the inhaler going off by accident.




When to switch to a new EASYHALER

The dose counter shows the number of remaining doses. The counter turns after every 5th click. When the dose counter starts turning red, there are 20 doses left.

If you do not already have a new EASYHALER, contact your doctor for a new prescription. When the counter reaches 0 (zero), you need to replace the EASYHALER.

If you use the protective cover, you can keep it and insert your new inhaler into it.



Remember

- 1. Shake – 2. Click – 3. Inhale.**
- After you have taken the dose, rinse your mouth with water and spit it out and/or brush your teeth.
- Do not get your inhaler wet, protect it from moisture.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

Revision Date:

June 2022



20-06-2023/MJ



Proof Colors:

■ Black

Size : 200 x 620 mm
Minimum Font 9 pt
Omnes Regular & Bold
Pharmacode No. 45