

BISLAN[®] TABLET

Ingredient(s):

Each tablet contains:

Bromhexine HCl 8mg

Pharmacodynamics:

Bromhexine HCl has a secreto-motor action without other side effects. Furthermore, it has a mucolytic action without the risk of poisoning. It is superior to its parent alkaloids, vasicine and vasicinone, which possess bronchial constriction and dilatation effects.

Pharmacokinetics:

- The increased bronchial fluid secretion caused by Bromhexine HCl is not inhibited by atropine. There is neither stimulation of vagus nerve nor are there tolerance or accumulation ever found.
- Bislan[®] Tablet's action sets in about twenty minutes after administration, reaching peak in 6~8 hours, hence it has the advantages of quickness in action and length of duration.
- Bromhexine HCl is well absorbed from the gastrointestinal tract. Its peak plasma concentration occurs after 1 hour of administration. Most of the dose is excreted in the urine. Ambroxol is a metabolite of bromhexine.

Indication(s):

A mucolytic agent used in the treatment of respiratory disorders associated with viscid or excessive mucus.

Dosage and Administration:

Adults : 1 tablet, three times daily.

Children : 5 - 10 years of age, 1/2 tablet, four times daily.

Under 5 years of age, 1/2 tablet, twice daily.

To be used as directed by a physician.

Route of administration:

Oral administration

Contraindication(s):

Contraindicated in patients known to have hypersensitivity to this product.

Precaution(s) / Warning(s):

Bromhexine HCl should be given cautiously to patients with gastric ulceration since mucolytics may disrupt the gastric mucosal barrier.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Side Effect(s) / Adverse Reaction(s):

Gastrointestinal : Nausea, anorexia, abdominal discomfort and abdominal pain may occur.

CNS : Bromhexine may occasionally cause headache.

Dermatological : Uncomfortable or allergic reactions (skin rash, pruritus, etc.), should these occur, discontinue use of this product.

Immune System Disorders : Frequency not known. Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders : Frequency not known. Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP)).

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdose. Management of the patient consists of symptomatic and supportive therapy.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product Description and Packing(s):

A white color tablet, one side impressed with a cross mark throughout the surface.

Plastic bottle of 1000's and 1500's (for export only).

Blister packing of 10's x 10 and 10's x 100.

Item Code	: 204140 - 11
PM Code	: Ins.P BLT
Measurement(mm) ± 2mm	: 260(L) x 110(W)
Cursor (Attachment 1)	: R 048 - 050 mm
Cursor (Attachment 2)	: B 018 - 022 mm
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

L
260mm (±2mm)

W
110mm (±2mm)

BISLAN[®] TABLET

Bahan Aktif:

Setiap tablet mengandungi:

Bromhexine HCl 8mg

Indikasi:

Sebagai agen mukolitik untuk merawat keadaan berkaitan dengan mukus yang berlebihan dan pekat.

Dos:

Dewasa : 1 tablet, 3 kali sehari.

Kanak-kanak : 5 - 10 tahun, 1/2 tablet, 4 kali sehari.

Bawah 5 tahun, 1/2 tablet, 2 kali sehari.

祛痰錠

成份：
每錠含：

Bromhexine HCl 8mg

適應症：

祛痰。

用法、用量：

成人 : 每次 1 錠，每日 3 次。

兒童 5 - 10 歲者 : 每次 1/2 錠，每日 4 次。

兒童 5 歲以下者 : 每次 1/2 錠，每日 2 次。

Item Code	: 204140 - 11
PM Code	: Ins,P BLT
Measurement(mm) ± 2mm	: 260(L) x 110(W)
Cursor (Attachment 1)	: R 048 - 050 mm
Cursor (Attachment 2)	: B 018 - 022 mm
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

L
260mm (±2mm)



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD., (199001001034)
 Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
 Kawasan Perindustrian Bandar Baru Bangi,
 43000 Kajang, Selangor, Malaysia.
 Ordering Line: 1 800 88 3027
 Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 18 Dec 2020

W
110mm (±2mm)