



DEXCHLORPHENIRAMINE FORTE SYRUP

COMPOSITION:

Each 5ml contains
Dexchlorpheniramine Maleate 2mg

Methylparaben 0.1% w/v and Sodium Benzoate
0.1% w/v as preservatives.
Ponceau 4R as colourant.

PRESENTATION:

A red coloured and strawberry flavoured syrup.

DOSAGE AND ADMINISTRATION:

For oral administration only.
Adult: 2.5ml to 5ml (1 to 2mg), 4 - 6 hourly, up to
a maximum of 30ml (12mg) / 24 hours.
Children 6 - 12 years: 2.5ml (1mg), 4 - 6 hourly,
up to a maximum of 15ml (6mg) / 24 hours.

INDICATIONS:

For allergic conditions including hay fever,
urticaria, angioedema, vasomotor rhinitis,
allergic eczema, atopic and contact dermatitis,
drug and serum reactions, insect bites, pruritus.

PHARMACOLOGY:*Mechanism of Action:*

Dexchlorpheniramine Maleate is an alkylamine
derivative with antihistamine and anticholinergic
effects. The antihistamine acts by competing
with histamine for H1 receptor sites on the
effector cells, to the exclusion of agonist
molecules, without itself initiating a response.
This antagonism is competitive and reversible. It
does not prevent the production of histamines.

Pharmacokinetics:

Dexchlorpheniramine Maleate is readily
absorbed from the gastrointestinal tract and
parenteral sites of administration. Following oral
administration, the effects may start within 15 to
30 minutes, reaching peak within 1 hour, and
may last about 4 - 6 hours. It is extensively
metabolised in the liver, and excreted mainly as
metabolites in the urine.

CONTRAINDICATIONS:

Contraindicated in patients with known
hypersensitivity to antihistamines.

PRECAUTIONS:

May cause drowsiness and hence may impair
ability to drive vehicles or operate machinery.
MAO inhibitors may prolong some of its actions.

Patients should be warned that antihistamines may enhance
the sedative effects of CNS depressants including alcohol,
barbiturates, hypnotics, narcotic analgesics, sedatives and
tranquilizers. Use with caution in patients with narrow angle
glaucoma, history of bronchial asthma, hyperthyroidism,
cardiovascular disease and prostatic hypertrophy. Use is not
recommended in nursing mothers, pregnancy, newborn and
premature infants. Safety and efficacy of the drug in children
younger than 2 years of age has not been established.

SIDE EFFECTS:

The most common side effect is sedation, varying from
drowsiness to deep sleep, and including inability to
concentrate, lassitude, dizziness, hypotension, muscular
weakness and incoordination. Sedative effects, when they
occur, may diminish after a few days. Other side effects
include gastrointestinal disturbances, such as nausea,
vomiting, diarrhoea or constipation, colic, and epigastric
pain. It may also produce headache, blurred vision, tinnitus,
elation or depression, irritability, nightmares, anorexia,
difficulty in micturition, dryness of the mouth, tightness of the
chest, tingling, heaviness and weakness of the hands. It may
potentiate CNS depressants. Large doses may precipitate
fits in epileptics.

DRUG INTERACTIONS:

MAO inhibitors may enhance the antimuscarinic effects of
dexchlorpheniramine maleate while CNS depressants
including alcohol, barbiturates, hypnotics, narcotic
analgesics, sedatives and tranquilizers may enhance
sedative effects. It is also important to note that
dexchlorpheniramine may mask the warning signs of
damage caused by ototoxic drugs such as aminoglycoside
antibiotics.

OVERDOSAGE AND TREATMENT :

If the drug has been taken recently by mouth, the stomach
should be emptied by aspiration and lavage. Emetics are
generally of little value. The patient should be kept quiet to
minimise the excitation which occurs particularly in children.
Convulsions and marked CNS stimulation should preferably
be treated with diazepam or phenobarbitone intramuscularly,
paraldehyde, thiopentone sodium, and chlorpromazine have
also been recommended. Severe hypotension may require
fluid replacement, and assisted respiration may be
necessary. Forced diuresis is of little value since
antihistamines are rapidly metabolised and only traces are
recovered in the urine.

STORAGE:

Keep container well close. Store below 30°C. Protect from
light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK****PACK QUANTITIES:**

Available in PET bottle of 60ml & 100ml.

Further information can be obtained from pharmacist,
physician or the manufacturer.

MAL19984387AZ

Product Registration Holder & Manufactured By:

 **Kotra Pharma (M) Sdn. Bhd.**
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision : 18/11/22