

YSP ANTIFUNGAL VAGINAL TABLET 100MG

Ingredient(s):

Each tablet contains:

Miconazole Nitrate 100mg

Pharmacodynamic:

Miconazole nitrate has a fungistatic effect resulting from its inhibition of membrane sterol synthesis and a fungicidal effect from impairment of the barrier function of the membrane.

Miconazole nitrate inhibits the fungal biosynthesis of triglycerides and phospholipids.

It also inhibits oxidative and peroxidative enzyme activity, resulting in intracellular accumulation of hydrogen peroxide which is toxic to the fungal cell.

In vitro studies on miconazole nitrate show that it possesses fungicidal action against *Candida albicans* and other species of this genus.

Miconazole has also demonstrated clinical efficacy against *Coccidioides immitis*, *Candida albicans*, *Cryptococcus neoformans*, *Petrellidium boydii*, and *Paracoccidioides brasiliensis*.

Pharmacokinetic:

Miconazole is widely distributed in body tissues; it penetrates well into inflamed joints, the vitreous humor of the eye, and the peritoneal cavity, but poorly into the saliva and sputum. It crosses the blood-brain barrier, but only to a small extent.

Miconazole is 90 % bound to serum proteins, it undergoes hepatic transformation and has a terminal elimination half-life of approximately 24.1 hours.

Approximately 14 to 22 % of the dose is excreted in the urine mainly as inactive metabolites.

Route of Administration:

Vaginal.

Indication(s):

For the local treatment of vulvovaginal candidiasis (moniliasis).

Dosage and Administration:

One tablet once daily. Insert into vagina before going to bed at night.

One therapeutic duration is 7 days.

To be used according to a physician's prescription.

Contraindication(s):

Patients known to be hypersensitive to this medication.

Warnings and Special Precautions:

1. If a reaction suggesting sensitivity or chemical irritation should occur, use of this medication should be discontinued.
2. For vaginal use only.
3. In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions)

Interaction with Other Medicaments

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and anticoagulant effect should be monitored (see Warnings and Special Precautions)

Pregnancy and Lactation:

None has been mentioned.

Side Effect(s) / Adverse Reaction(s):

Most complaints are vulvovaginal burning, itching or irritation. Other complaints such as vaginal burning, pelvic cramps, hives, skin rash and headache occur rarely.

Symptoms and Treatment for Overdosage, and Antidote(s):

Overdosage of Miconazole in humans has not been reported to date. Should overdosage occurs, management of the patient may consist symptomatic and supportive therapy.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):

A white color long shaped tablet, one side impressed with a mark '7118', and the other side with wording of 'Yung Shin'.

Plastic bottle of 100's, 500's and 1000's. (For export only)

Strip packing of 10's x 10.

Please consult your physician/pharmacist before using this product if you are on the anticoagulant medicine warfarin, because bleeding from the nose/gums or bruising may occur spontaneously.

Sila dapatkan nasihat doktor atau ahli farmasi sebelum menggunakan keluaran ini jika anda mengambil ubat warfarin, iaitu sejenis ubat antipembekuan darah, kerana lebam/pendarahan pada gusi/hidung boleh berlaku secara spontan.

YSP ANTIFUNGAL VAGINAL TABLET 100MG

Bahan Aktif:

Setiap tablet mengandungi:

Miconazole Nitrate 100mg

Indikasi:

Untuk rawatan setempat bagi kandidiasis vulvovaginal (moniliasis).

Dos:

1 tablet sehari. Masukkan ke dalam vaginal sebelum waktu tidur.

Tempoh untuk satu rawatan lengkap adalah selama 7 hari.

Digunakan seperti yang diarahkan oleh doktor.

無 黴 陰道錠

成份:

每錠中含:

Miconazole Nitrate 100mg

適應症:

對黴菌、酵母菌、細菌之單獨或混合感染均有效。針對念珠菌性陰道炎、感染性白帶及其他具有感受性病原體所引起之感染症。

用法 用量:

每日1次，於就寢前以1粒置入陰道深處，連續7天為一治療程。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679