
DUOGYNON INJECTION (1 ML AMP)

DESCRIPTION: DUOGYNON INJECTION (1 ML AMP): A clear, slightly yellow solution

COMPOSITION: Each 1 ml of injection contains Progesterone 50 mg and Estradiol Benzoate 3 mg

PHARMACODYNAMICS: DUOGYNON acts in the endometrium by converting the proliferative phase induced by oestrogen to a secretory phase and preparing the uterus to receive the fertilized ovum.

PHARMACOKINETICS: The progesterone in DUOGYNON Injection is rapidly absorbed from the site of the intramuscular injection. The half-life in blood is only a few minutes. It is metabolized in the liver about 12% to pregnanediol which is excreted in the urine conjugated with glucuronic acid. Progesterones are excreted in the milk of nursing mothers. The oestradiol is partly bound to plasma proteins and is rapidly metabolized mainly in the liver and excreted in the urine as sulphate and glucuronide esters together with a small proportion of unchanged estradiol. Oestrogens are also excreted in the milk of nursing mothers.

INDICATION: Secondary amenorrhoea of short duration

RECOMMENDED DOSAGE: Pregnancy must be excluded before the administration of DUOGYNON Injection. One single injection of DUOGYNON 1 ml (Progesterone 50 mg and Estradiol Benzoate 3 mg) usually induces a menstruation like withdrawal bleeding within 3-6 days; in isolated cases late. Further diagnostic and therapeutic measures are required if no bleeding occurs within 2 weeks.

As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates
--

ROUTE OF ADMINISTRATION: For intramuscular injection

CONTRAINDICATIONS: Contraindicated in pregnancy. A history of herpes of pregnancy. Thrombophlebitis, thromboembolic disorders, CVA or history of these conditions; liver disease or history of cholestatic jaundice, carcinoma of breast or genital organs, suspected estrogen-dependent neoplasia; disturbance of lipometabolism; undiagnosed vaginal bleeding, hemiplegic migraine.

WARNINGS & PRECAUTIONS: Avoid giving to diabetes mellitus patients if either retinopathy or nephropathy is present. Avoid patients with family history of arterial disease (ischaemic heart disease, ischaemic cerebrovascular accident). History of depression; uterine fibroids; lactation, epilepsy and migraine.

INTERACTION WITH OTHER MEDICAMENTS: Effectiveness may be reduced by certain antibiotics, phenytoin, carbamazepine, barbiturates, rifampicin.

USE IN PREGNANCY & LACTATION: Contraindicated in pregnancy. A history of herpes of pregnancy.

SIDE EFFECTS: Nausea, vomiting, headache, impairment of liver function, reduced menstrual loss, "spotting" in early cycles.

SYMPTOMS AND TREATMENT OF OVERDOSE: This is unlikely and is not expected to produce any adverse effects. Treatment is observation and, if necessary, symptomatic and supportive measures should be provided.

INCOMPATIBILITIES: None known

STORAGE CONDITIONS:
Store below 30°C. Protect from light. Keep out of reach of children. *Jauhkan daripada kanak-kanak.*

SHELF LIFE:
Please refer to outer package.

PACK SIZE: Packs of 10 x 1 ml ampoule

PRODUCT REGISTRATION HOLDER & MANUFACTURER:
DUOPHARMA (M) SDN BHD
Lot 2599, Jalan Seruling 59 Kawasan 3,
Taman Klang Jaya, 41200 Klang, Selangor, MALAYSIA.