

11 cm

PREDON

LF-0038

Composition :
Each mL contains : Prednisolone Acetate 25 mg

Preservatives Used :
Methyl paraben : 1.65 mg
Propyl paraben : 0.18 mg

Product Description : Sterile white suspension for injection.

Pharmacodynamics :
Prednisolone is used especially as anti-inflammatory in the treatment of conditions where a cortisone-like action is needed. It is usually preferred to cortisone in such conditions as rheumatoid arthritis, rheumatic fever, status asthmaticus and ulcerative colitis. It is useful in the treatment of nephrotic edema, as it produces diuresis. It is not used alone in the treatment of adrenal deficiency state because of its weak salt-retaining effect.

Pharmacokinetics :
Absorption : Prednisolone is absorbed from the gastrointestinal tract and readily converted to prednisolone in the liver and has a biological half-life of about 60 minutes.
Distribution : Protein binding, about 70% - 90% bound to plasma protein.
Metabolic reaction : Reduce to 20 beta-dihydro-prednisolone hydrocortisone and conjugation occur to some extent.
Excretion : About 10-30 % of a dose is excreted in the urine as metabolite, 60% - 80% of which is conjugated, a dose is almost completely excreted in 3 days. Prednisolone is excreted in milk in insignificant amounts.

Indication : Rheumatoid arthritis, inflammatory and allergic conditions, bronchial asthma.

Recommended Dose :
Intramuscular : 2 to 30 mg, 2 times a day.
Intra-articular : 12.5 to 25 mg at each site every one to three weeks

Route of Administration : Intra-muscular (IM) & Intra-articular (IA)

Contraindications :
It is contraindicated to patients in systemic fungal infections, pregnancy, lactation and in those persons who have shown hypersensitivity to any component of this preparation. Patients should not undergo immunizations. It is contraindicated to patients with tuberculosis, except as adjunctive treatment with tuberculostatic drugs.

Warnings and Precautions :
Extreme caution in congestive heart failure, diabetes mellitus, chronic renal failure, hypertension, recent intestinal anastomoses. Infectious diseases, uraemia, elderly person, diverticulitis, myasthenia gravis and ocular herpes simplex (corneal perforation may occur). History of peptic ulcer or psychotic tendencies; family history of diabetes mellitus; patients with previous tubercular infection should receive hemoprophylaxis if corticosteroid therapy is prolonged. Monitor spinal radiographs for signs of osteoporosis; rheumatoid arthritis patients and postmenopausal women are at special risk. Growth inhibition or arrest, which cannot be overcome with exogenous growth hormone, may occur in children given relatively low doses. Prolonged therapy may produce posterior subcapsular cataracts, Glaucoma with possible damage to optic nerve, steroid myopathy and Cushing's habitus may develop with long-term therapy. Hepatic cirrhosis and hypothyroidism may enhance corticosteroid effects.
Not for IV use.
This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.
Pheochromocytoma crisis, which may be fatal, has been reported after administration of systemic corticosteroids. Corticosteroids should only be administered to patients with suspected or identified pheochromocytoma after an appropriate risk/benefit evaluation.
Scleroderma renal crisis
Caution is required in patients with systemic sclerosis because of an increased incidence of (possibly fatal) scleroderma renal crisis with hypertension and decreased urinary output observed with a daily dose of 15 mg or more prednisolone.

Interactions with other Medicaments :
Response to corticosteroids is reduced by concurrent administration of barbiturates, phenylbutazone, phenytoin or rifampicin. Response to anti coagulants and salicylates is reduced by corticosteroids. Abrupt cessation of prolonged corticosteroid therapy produces acute adrenal insufficiency which may be life-threatening.

Pregnancy and Lactation :
Pregnancy category : B
Animal studies : Birth defects reported in mice, rats and rabbits.
Human studies : Adequate studies of pregnant women are not available.
Avoid completely during the first 3 months. Limit use during the final 6 months as much as possible. If used, examine infant for possible deficiency of adrenal gland function.
Lactation : Presence of this drug in breast milk is found but amounts received are generally less than 0.1% of a therapeutic dose. Consult the physician.

Side Effects :
In the course of corticosteroid treatment, especially when intensive and prolonged the following adverse effects may occur :
- Alterations in the water-electrolyte balance which in particularly predisposed patients may cause hypertension or congestive heart failure.
- Alterations in the musculoskeletal system such as osteoporosis, myopathies, brittleness of bones.
- Alterations in the gastrointestinal tract with possible peptic ulcer, or recurrence of peptic ulcer.
- Skin changes : thinning and fragility of the skin, delayed scar formation.
- Neurological disorders : dizziness, headache, increased intracranial pressure.
- Endocrine disorders : such as menstrual irregularities, Cushing-like syndromes, delayed growth in children, inhibition of the pituitary adrenal axis, reduced tolerability to glucose with possible appearance of a latent diabetes mellitus.
- Ophthalmic complications such as posterior subcapsular cataract, increased intraocular pressure, glaucoma exophthalmos.
- Increased protein catabolism resulting in negative nitrogen balance.

Symptoms and Treatment of Overdose :
Symptoms : Anxiety, depression and or stimulation, stomach bleeding, increased blood sugar, high blood pressure and retention of fluid.
Treatment : The patient should be taken to the hospital immediately for stomach pumping oxygen, intravenous fluid and other symptomatic treatment.

Effects on Ability to Drive and Use Machine : This medicine has no or negligible influence on the ability to drive and use machines.

Storage Condition : Store below 30 °C, Protect from light.

ATC Code : H02AB06 - Prednisolone

MAL Number : MAL19986420AZ

Shelf Life : 4 years

Pack Size : 1 vial of 10 mL

Date of Revision : 21 July 2022

Manufactured by :
 **L.B.S. LABORATORY LTD., PART.**
602 Soi Panichnant, Sukhumvit 71 Rd., Bangkok 10110 Thailand,
Tel : (66) 2392-4082 Fax : (66) 2391-7122

Product Registration Holder :
HEALOL PHARMACEUTICALS SDN. BHD.
74-3, Jalan Wangsa Delima 6, KLSG Wangsa Maju, 53300 Kuala Lumpur, Malaysia.
Tel : (60) 3-4142-7106, 4142-7107 Fax : (60) 3-4146-7108

15 cm

Code: R 114/22
Project: PREDON
LF E
Date: 21-07-2022
Printed Size: 11 x 15 cm (WxH)
Print Colours:
Process Colours: CMYK **K=100**
Special Colours:

R114-22_PREDON_LF E_ML_1stDraft_SM_21-07-2022