

YSP BISACODYL SUPPOSITORIES

Presentation(s):

Bisacodyl Suppositories 10mg

Each supp. contains:

Bisacodyl 10mg

Bisacodyl Suppositories 5mg

Each supp. contains:

Bisacodyl 5mg

Pharmacodynamics:

Bisacodyl is a stimulant or contact laxative. It increases the peristalsis of the colon by a direct effect on the smooth intestinal musculature by stimulation of intramural nerve plexi. It also has been shown to promote fluid and ion accumulation in the colon to increase the laxative effect. Bisacodyl suppository can be used in functional constipation to facilitate defecation. In preparation for surgery proctoscopy, or radiologic examination, bisacodyl provides satisfactory cleaning of the bowel, obviating the need for an enema.

Pharmacokinetics:

Bisacodyl acts within 30 minutes when given rectally. The contact action of the drug is restricted to the colon, and motility of the small intestine is not appreciably influenced. Local axon reflexes, as well as segmental reflexes are initiated in the region of contact and contribute to the widespread peristalsis activity producing evacuation.

For this reason Bisacodyl may often be employed satisfactorily in patients with ganglionic blockade or spinal cord damage (paraplegia, poliomyelitis, etc.).

Indication(s):

- i) Constipation
- ii) Bowel preparation for radiological procedures and surgery.

Dosage and Administration:**i) Constipation :**

Adults and children over 10 years: 1 suppository (10mg) for immediate effect daily.

Children 4 – 10 years: 1 suppository (5mg) for immediate effect daily.

ii) Bowel preparation:

Adults and children over 10 years: 1 suppository (10mg) on the following morning before procedure.

Children aged 4 -10 years of age: 1 suppository (5mg) on the following morning before procedure.

Bisacodyl suppository is usually effective within 30 minutes after insertion.

In the management of constipation, once regularity has been restarted dosage should be reduced and can usually be stopped.

Route of Administration:

For rectal use.

Contraindication(s):

Patients with intestinal obstruction, acute surgical abdominal conditions such as acute appendicitis, acute inflammatory bowel disease or with undiagnosed abdominal symptoms.

Side Effect(s) / Adverse Reaction(s):

As with any laxative, abdominal cramps are occasionally noted, particularly in severely constipated individuals. When administered rectally it sometimes cause local irritation.

Precaution(s) / Warning(s):

- 1. Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.
- 2. Care should be taken in patients with inflammatory bowel disease.
- 3. Should be given with caution in patients with rectal fissures or ulcerated haemorrhoids.
- 4. Prolonged use should be avoided.
- 5. As with all medicines, keep these suppositories out of reach of children.

Drug Interaction(s):

At high doses, risk of electrolytes imbalance increased with concurrent use of diuretics and adrenocorticoids.

Symptoms and Treatment for Overdosage, and Antidote(s):

Symptoms of acute overdosage include confusion, irregular heartbeat, abdominal muscle cramps, unusual tiredness or weakness.

There is no specific antidote for its overdose, therefore, management of the patients consist of symptomatic and supportive therapy.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 25°C. Protect from light and moisture.

Product Description and Packing (s):

Bisacodyl Suppositories 10mg

A white to off-white spindle suppository.

10 supp. per strip, 5 strips per box

10 supp. per strip, 10 strips per box.

Bisacodyl Suppositories 5mg

A white to off-white spindle suppository.

10 supp. per strip, 10 strips per box.

5 supp. per strip, 2 strips per case.

5 supp. per strip, 1 strip per case.

YSP BISACODYL SUPOSITORI

Bahan Aktif:

Bisacodyl Supositori 10mg

Setiap supp. mengandungi:

Bisacodyl 10mg

Bisacodyl Supositori 5mg

Setiap supp. mengandungi:

Bisacodyl 5mg

Indikasi:

Sembelit.

Dos:

Dewasa: 1 supositori, 10mg, sekali sehari. Kanak-kanak berumur 4 - 10 tahun: 1 supositori, 5mg, sekali sehari. Dimasukkan dengan sempurna ke dalam rektum. Supositori Bisacodyl biasanya berkesan dalam masa 30 minit.

YSP 無秘 栓劑

成份:

Bisacodyl Supositori 10mg

每粒中含:

Bisacodyl 10mg

Bisacodyl Supositori 5mg

每粒中含:

Bisacodyl 5mg

適應症:

便秘。

用法、用量:

通常成人每天1個10毫克，小孩4-10

歲：每天1個5毫克栓劑，塞入肛門深部。



Manufacturer and Product Registration Holder :
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679