

FRONT

ALUCID TABLET

For Antacid and Anti-flatulent

PRODUCT DESCRIPTION

A 16mm in diameter, round, flat, marked 'Alucid' on both side, yellow tablet

INDICATIONS

- **Alucid Tablet** can be used for relief of gastric pain, hyperacidity, flatulence and heartburn.
- **Alucid Tablet** will relieve and control pain in the symptomatic management of gastric and duodenal ulcers and reflux oesophagitis by neutralising hydrochloric acid in the gastric secretion.
- **Alucid Tablet** is also suitable as a domestic remedy in conditions affecting the stomach which may not necessarily be related to hyperacidity.
- **Alucid Tablet** comes with a cooling and refreshing peppermint flavour.

CONTENTS

Each Tablet of **Alucid Tablet** contains:-
Aluminium Hydroxide Dried Gel 250mg
Magnesium Hydroxide 250mg
Simethicone 50mg

DOSAGE

Adult and children above 12 years : 1 to 2 tablets to be chewed before meals or as directed by physician.

Route of administration: Oral

SIDE EFFECTS

Constipation, diarrhoea, intestinal obstruction might happen if a larger than recommended dose of **Alucid Tablet** been taken.

PHARMACODYNAMICS

These medications react chemically to neutralize or buffer existing quantities of stomach acid but have no direct effect on its output. This action results in increased pH values of stomach contents, thus providing relief of hyperacidity symptoms. Also, these medications reduce acid concentration within the lumen of the esophagus. This causes an increase in intra-esophageal pH and a decrease in pepsin activity. Antacids may increase lower esophageal sphincter (LES) pressure. Aluminium containing antacids have a cytoprotective effect on the gastric mucosa that may be associated with stimulation of prostaglandin secretions, thus providing protection against mucosal necrosis and hemorrhage caused by corrosive agents, such as aspirin and ethanol.

PHARMACOKINETICS

Aluminium containing: small amounts of the aluminium in aluminium hydroxide are absorbed from the intestine. Magnesium containing: approximately 10% of the magnesium in magnesium hydroxide is absorbed from the intestine. Simethicone: Acts in vitro to lower the surface tension of gas bubble. When administered by mouth causes bubbles of gas in the gastro-intestinal tract to coalesce, thus aiding their dispersion.

CONTRAINDICATION

Alucid Tablet should have not be taken by person who suffers from severe renal function impairment.

PRECAUTIONS

Use with care in renal insufficiency. Should not be used if symptoms of appendicitis are present. Do not take within 1 or 2 hours of other oral medication as it may impair their absorption. Please consult your physician if pain persists.

This product contains aspartame and is not suitable for phenylketourics.

PACKING

18 x 10's.

Not all pack sizes are available locally.

STORAGE

- STORE IN A DRY PLACE BELOW 30°C
- PROTECT FROM LIGHT
- KEEP OUT OF REACH OF CHILDREN

SHELF LIFE

Please refer to box.

DRUG INTERACTION:

Because aluminium and magnesium salts reduce the gastrointestinal absorption of tetracyclines, the use of Alucid Tablet should be avoided during oral tetracycline therapy.

The use of antacids containing aluminium can reduce the absorption of drugs, in particular H2-antagonists, atenolol, cefdinir, cefpodoxime, chloroquine, tetracyclines, dasatinib monohydrate, diflunisal, digoxin, dexamethasone, bisphosphonates, eltrombopag, elamine, elvitegravir, ethambutol, fluoroquinolones, glucocorticoids, indomethacin, iron salts, isoniazide, ketoconazole, levothyroxine, lincosamides, metoprolol, nilotinib, phenothiazine neuroleptics, penicillamines, propranolol, raltegravir, potassium, rilpivirine, ricitquat, rosuvastatin, sodium fluoride, and antiviral treatments in combination with tenofovir alafenamide fumarate/emtricitabine/sodium bictegravir.

- Polystyrene sulfonate (Kayexalate)

Caution is advised when this medicinal product is used concomitantly with polystyrene sulfonate (Kayexalate) due to the potential risk of reduced effectiveness of the resin in binding potassium, of metabolic alkalosis in patients with renal impairment (reported with aluminium hydroxide and magnesium hydroxide) and of intestinal obstruction (reported with aluminium hydroxide).

- Aluminium hydroxide and citrates may cause hyperalbuminemia, especially in patients with renal impairment.

As magnesium hydroxide administration results in urine alkalization, increased excretion of salicylates has been observed in the event of concomitant use.

As a precaution, allow at least 2 hours (4 for fluoroquinolones) to elapse between taking any oral drug and Alucid Tablet.

Concomitant use of aluminium hydroxide and citrates may result in increased aluminium levels, especially in patients with renal impairment. Urine alkalization following magnesium hydroxide administration may affect the excretion of some drugs; increased excretion of salicylates has been observed.

PREGNANCY AND LACTATION:

There are no available data on Alucid Tablet use in pregnant women. No conclusions can be drawn regarding whether or not Alucid Tablet is safe for use during pregnancy.

Breast-feeding

The medicinal product should only be used if absolutely necessary, under direct medical supervision, after having evaluated the expected benefits for the mother compared with the possible risk to the fetus or infant.

Breast-feeding

Because of the limited maternal absorption when taken in accordance with the indicated dosage regimen, aluminium hydroxide and its magnesium salt combinations are considered compatible with breast-feeding.

Effects on ability to drive and use machines:

Alucid Tablet has no influence on the ability to drive and use machines

OVERDOSE:

There is very limited experience with deliberate overdose. Cases of overdose with aluminium salts are more likely to occur in patients with severe chronic renal impairment, with the following symptoms: encephalopathy, seizures and dementia, hypermagnesemia.

The most frequently reported symptoms of acute overdose with aluminium hydroxide and in combination with magnesium salts include diarrhea, abdominal pain and vomiting.

Large doses of this medicinal product may cause or aggravate intestinal obstruction and ileus in patients at risk.

Treatment should be symptomatic and include general supportive measures.

Aluminum and magnesium are eliminated through the urinary route; treatment of magnesium overdose consists of rehydration and forced diuresis. In case of renal function deficiency, hemodialysis or peritoneal dialysis is necessary.

ALUCID 药品

缓和胃酸及气胀

说明

- **Alucid 药品**有助于减轻胃痛、胃酸过多、气胀和胃灼热。

- **Alucid 药品**有助于中和胃部的氢氟化酸，进而控制及减轻胃和十二指肠溃疡造成的疼痛以及食道逆流症状。

- **Alucid 药品**也适于治疗不是因胃酸过多引起的胃痛。

- **Alucid 药品**备有清爽冰凉的薄荷味。

成份

每1颗Alucid 药品含有：
250 毫克 Aluminium Hydroxide Dried Gel
250 毫克 Magnesium Hydroxide
50 毫克 Simethicone

剂量

成人及12岁以上儿童：在饭前嚼嚼1至2片或遵照医生指示。

副作用

若服用大量**Alucid 药品**，可能出现便秘、腹泻及肠胃阻塞现象。

禁忌

患有严重肾脏功能衰竭者请勿服用**Alucid 药品**。

警告

肾脏机能不全者应谨慎服用。如果出现阑尾炎症状，请勿服用。请勿服用其他口服药物之后的1至2小时内服用**Alucid 药品**，因为此举可能会削弱其吸收力。

如果疼痛持续，请询问您的医生。

本产品含有阿斯巴甜和不适合苯丙酮尿症。

包装

18 x 10's

贮藏

- 置于乾燥及摄氏30°C 以下。
- 远离光线。
- 置于孩童无法触及之处。

贮藏期

请参阅盒内中文说明

size = 155 x 178mm

BACK

ALUCID TABLET

Sediaan Antasid dan Anti-flatulen

INDIKASI

- **Alucid Tablet** digunakan untuk melegakan kesakitan akibat gastrik, asid berlebihan, flatulen serta pedih di bahagian hulu hati.
- **Alucid Tablet** digunakan untuk melegakan dan mengawal kesakitan akibat gastrik dan ulser di bahagian bawah dan atas usus perut dengan meneutralkan asid hidroklorik di bahagian berkenaan.
- **Alucid Tablet** juga sesuai digunakan sebagai rawatan kepada sakit perut yang tidak berkaitan dengan kandungan asid yang berlebihan.
- **Alucid Tablet** mempunyai rasa pudina yang nyaman dan menyegarkan.

KANDUNGAN

Setiap **Alucid Tablet** mengandungi :-

Aluminium Hydroxide Dried Gel 250mg

Magnesium Hydroxide 250mg

Simethicone 50mg

SUKATAN

Dewasa dan kanak-kanak 12 tahun ke atas:

1 hingga 2 biji tablet dikunyah sebelum makan atau seperti arahan oleh pakar perubatan.

KESAN SAMPINGAN

Sembelit, cirit-birit dan jangkitan usus tersumbat mungkin berlaku jika dos **Alucid Tablet** diambil secara berlebihan.

PERHATIAN

Alucid Tablet tidak boleh digunakan oleh pesakit buah pinggang yang serius.

AMARAN

Gunakan secara berhati-hati sekiranya anda mengalami masalah buah pinggang. Jangan gunakan sekiranya anda mempunyai tanda-tanda penyakit appendik. Tidak boleh diambil 1 atau 2 jam selepas mengambil ubat-ubatan lain kerana ia mungkin memberi kesan terhadap keberkesananannya.

Sila rujuk kepada pakar perubatan sekiranya sakit berlarutan.

Produk ini mengandungi aspartame dan tidak sesuai untuk phenylketourics.

BUNGKUSAN

18 x 10's

PENYIMPANAN

- SIMPAN DI TEMPAT KERING DI BAWAH SUHU 30°C.
- JAUHKAN DARI CAHAYA.
- JAUHI DARIPADA KANAK-KANAK.

JANGKA HAYAT

Sila rujuk pada kotak.

INTERAKSI UBAT:

Oleh kerana garam aluminium dan magnesium mengurangkan penyerapan dalam saluran gastrousus, penggunaan Alucid Tablet harus dielakkan semasa terapi tetrasiklin secara oral. Penggunaan antasid yang mengandungi aluminium boleh mengurangkan penyerapan ubat-ubatan tertentu, terutamanya antagonis H₂, atenolol, cefdinir, cefpodoxime, chloroquine, tetracyclines, dasatinib monohydrate, diflunisal, digoxin, dexamethasone, bisphosphonates, eltrombopag olamine, elvitegravir, ethambutol, fluoroquinolones, glucocorticoids, indomethacin, garam besi, isoniazide, ketoconazole, levothyroxine, lincosamides, metoprolol, nilotinib, phenothiazine neuroleptics, penicillamines, propranolol, raltegravir potassium, rilpivirine, riociguat, rosuvastatin, sodium fluoride, rawatan antiviral bersama tenofovir tenofovir alafenamide fumarate/emtricitabine/sodium bictegravir.

- Polistirena sulfonat (Kayexalate)

Berhati-hati apabila produk ubat ini digunakan bersama dengan polistirena sulfonat (Kayexalate) kerana terdapat potensi risiko pengurangan keberkesanan resin dalam mengikat kalium, risiko alkalosis metabolik pada pesakit yang mengalami masalah kegagalan buah pinggang (dilaporkan dengan aluminium hidroksida dan magnesium hidroksida) dan halangan usus (dilaporkan dengan aluminium hidroksida).

- Aluminium hidroksida dan sitrat boleh menyebabkan hiperalbuminemia, terutamanya pada pesakit yang mengalami masalah kegagalan buah pinggang. Memandangkan pemberian magnesium hidroksida mengakibatkan pengalkalian air kencing, peningkatan perkumuhan salisilat telah diperhatikan apabila digunakan bersama.

Sebagai langkah berjaga-jaga, pastikan terdapat selang masa sekurang-kurangnya 2 jam (4 untuk fluoroquinolones) antara pengambilan sebarang ubat oral dan Alucid Tablet.

Penggunaan serentak aluminium hidroksida dan sitrat boleh mengakibatkan peningkatan paras aluminium, terutamanya pada pesakit yang mengalami masalah kegagalan buah pinggang. Pengalkalian air kencing selepas pengambilan magnesium hidroksida boleh menjejaskan perkumuhan sesetengah ubat; peningkatan perkumuhan salisilat telah diperhatikan.

KEHAMILAN DAN PENYUSUAN:

Tiada data tersedia mengenai penggunaan Alucid Tablet pada wanita hamil. Tiada kesimpulan yang boleh dibuat sama ada Alucid Tablet selamat untuk digunakan semasa kehamilan atau tidak.

Kehamilan

Produk ubat ini hanya boleh digunakan jika benar-benar perlu, di bawah pengawasan perubatan langsung, selepas menilai manfaat yang dijangkakan untuk ibu berbanding dengan risiko yang mungkin berlaku kepada janin atau bayi.

Penyusuan susu ibu

Disebabkan penyerapan ubat oleh ibu adalah terhad apabila diambil mengikut rejimen dos yang disyorkan, kombinasi aluminium hidroksida dan garam magnesium dianggap sesuai dan serasi dengan penyusuan.

Kesan terhadap keupayaan untuk memandu dan mengendalikan mesin:

Alucid Tablet tidak memberi sebarang kesan terhadap keupayaan untuk memandu dan mengendalikan mesin.

TERLEBIH DOS:

Pengalaman mengenai kes terlebih dos secara sengaja adalah sangat terhad. Kes terlebih dos dengan garam aluminium lebih berkemungkinan berlaku di kalangan pesakit yang mengalami kegagalan buah pinggang kronik yang teruk, dengan simptom berikut: ensefalopati, sawan dan demensia, hipermagnesemia.

Simpom terlebih dos akut yang paling kerap dilaporkan dengan aluminium hidroksida dan gabungan garam magnesium termasuk cirit-birit, sakit perut dan muntah.

Pengambilan dos besar produk ubat ini boleh menyebabkan atau memburukkan lagi halangan usus dan ileus pada pesakit yang berisiko.

Rawatan harus berdasarkan simptom yang ditunjukkan dan merangkumi langkah-langkah sokongan umum.

Aluminium dan magnesium disingkirkan melalui laluan air kencing; rawatan untuk terlebih dos magnesium terdiri daripada rehidrasi dan diuresis paksa. Sekiranya terdapat kekurangan fungsi buah pinggang, hemodialisis atau dialisis peritoneal diperlukan.

Product Registration Holder:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Revision date: Sep 2025

1500012280 N.3

