
REXOM™ WINA ANTISEPTIC LIQUID

Composition and Strength of Active Ingredients:

Chlorhexidine Gluconate	0.3% w/v
Cetrimide	3.0% w/v

Description:

Product Description

Clear, orange solution

Pharmacodynamics

Chlorhexidine is an effective antiseptic with a wide range of activity against microorganisms, including gram positive and gram negative bacteria, fungi and viruses.

Cetrimide is a quaternary ammonium compound with surfactant and antiseptic properties.

Pharmacokinetics

Not applicable.

Indications

It is indicated as antiseptic for minor burns, cuts, wounds, sores and rashes.

It is used as general disinfectant, household disinfectant and personal hygiene.

Recommended Dose

Personal hygiene, wounds & burns: dilute 1 capful (10 ml) in 125 ml of water.

House disinfectant: dilute 1 capful (10 ml) in 250 ml of water.

Route of Administration

For topical application.

Contraindications

Hypersensitivity to chlorhexidine gluconate and cetrimide may occur after prolonged and repeated applications to the skin.

Warning and Precautions

Rexom Wina Antiseptic Liquid contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. Rexom Wina Antiseptic Liquid should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Keep all medicines out of the reach of children.

Keep out of the eyes and ears.

Treat accidental splashes by washing open eyes with water for at least ten minutes, seek medical advice.

Interactions with Other Medicaments

None known.

Pregnancy & Lactation

No known restriction

Side Effects

Immune system disorders

Frequency not known: Hypersensitivity including anaphylactic shock

Chemical burns in neonates (frequency unknown)

Symptoms and Treatment of Overdose

If swallowed, wash out of mouth and drink plenty of milk or water; seek medical advice and show the container to the doctor.

Effects on Ability to Drive and Use Machine

The ability to drive or use machines is not affected.

Storage Condition

Store below 30°C. Keep container tightly closed. Protect from light.

Presentation:

250ml and 350ml bottle

Registration Number:

MAL19991207X

Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

Manufactured by:

KCK Pharmaceutical Industries Sdn. Bhd.

No. 5 & 7 Hala Beringin,

Diamond Valley Industrial Park, 11960 Bayan Lepas,

Pulau Pinang, Malaysia.

Date of Revision:

04 November 2024