

TRETINON® CREAM 0.05%

Ingredient(s):

Each gram contains:

Tretinoin (Retinoic Acid)	0.5mg
Sorbic Acid (as preservative).....	2mg

Pharmacodynamics:

1. Tretinoin is a Vitamin A derivative used in the treatment of acne vulgaris. Although the exact mode of action of tretinoin is unknown, current evidence suggests that topical tretinoin appears to act by increasing epidermal cell mitosis and epidermal cell turnover, possibly resulting in the production of a less cohesive horny cell layer, which is more easily peeled off. This action facilitates the removal of existing comedones and may inhibit the formation of new comedones.
2. It has been proposed that increased turnover in the follicular epithelium prevents blocking by keratinous plugs.
3. Tretinoin has also been reported to suppress keratin synthesis.

Pharmacokinetics:

1. After topically applied tretinoin, some systemic absorption probably occurs, since small percentage of dose appears in urine; absorption may be increased when medication is applied to large surface areas, or for long periods of time in chronic extensive dermatoses, such as ichthyosis. About 5% of a topically applied dose of tretinoin is recovered in the urine.

Indication(s):

Topical treatment of acne vulgaris, in which comedones, papules and pustules predominate.

Dosage and Administration:

Apply once daily to the affected areas, preferably at bedtime, after cleansing with soap and water. Allow drying before applying medication.

Following application of the preparation, a short sensation of warmth and burning may be experienced. During the 1st to 5th week of treatment, slight reddening of the skin may occur. In cases of too severe skin irritation, the intervals between applications should be prolonged. Some redness and skin peeling may occur after application for several days but settles with time.

Therapeutic results may be noticeable after 2 to 3 weeks of therapy, but more than 6 weeks of therapy may be required for optimal results.

Route of administration:

Topical

Contraindication(s):

1. Patients who have demonstrated hypersensitivity to the drug.
2. Eczema, broken or sunburned skin.
3. Personal or family history of cutaneous epithelioma.
4. Pregnancy (see Section Pregnancy and Lactation)
5. Women planning a pregnancy

Precaution(s) / Warning(s):

1. For external use only. Excessive use should be avoided. Tretinoin should not be applied to the eyes, mouth or other mucous surfaces.
2. Exposure to excessive sunlight, including sunlamps, should be avoided while using the preparation, or a suitably effective sunscreen should be employed.
3. Cosmetics may be used, but the areas to be treated should be cleansed thoroughly before the medication is applied.
4. Concomitant use of other peeling medication may increase the irritant effect. Particular caution should be exercised in using preparations containing sulfur, resorcinol or salicylic acid with tretinoin.
5. If patient experiences severe or persistent irritation, the patient should be advised to discontinue application completely and if necessary consult a physician.
6. Children: Safety and effectiveness in children have not been established.

Pregnancy and Lactation:

Orally administered retinoids have been associated with congenital abnormalities. When used in accordance with the prescribing information, topically administered retinoids are generally assumed to result into low systemic exposure due to minimal dermal absorption. However, there could be individual factors (e.g. damaged skin barrier, excessive use) that contribute to an increased systemic exposure.

Tretinon Cream 0.05% is contraindicated (see Section Contraindications) in pregnancy, or in women planning a pregnancy.

If the product is used during pregnancy, or if the patient becomes pregnant while taking this drug, treatment should be discontinued.

Drug Interaction(s):

Special caution is indicated in case of simultaneous treatment with other preparations, especially desquamative agents such as sulfur, resorcinol and salicylic acid. Concomitant use may increase the effect of irritation.

Side Effect(s) / Adverse Reaction(s):

Local inflammatory reactions may occur, such as erythema, itching, irritation, stinging or burning, dryness, crusting, edema of the face, hyperpigmentation,

hypopigmentation, increase in the number of inflammatory lesions acne flare and excessive reaction to sun exposure.

All adverse reactions observed are reversible when treatment is discontinued.

Symptoms and Treatment for Overdosage, and Antidote(s):

If medication is applied excessively, no more rapid or better results will be obtained and marked redness, peeling, or discomfort may occur. Oral ingestion of the drug may lead to the same side effects as those associated with excessive oral intake of vitamin A. Reduce amount or frequency of application if undesirable reactions occur.

Storage Condition(s):

Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product Description and Packing(s):

Yellow cream.

Aluminium tube of 20g.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679