

Axcel® FLEMIN EXPECTORANT

COMPOSITION:

Each 5ml contains
Diphenhydramine HCl 12.5mg
Ammonium Cl 130mg

Sodium Benzoate 0.1% w/v as preservative &
Sodium Citrate 60mg/5ml.

PRESENTATION:

Brown colour syrup with caramel flavour.

INDICATIONS:

Relief of cough & alleviate nasal & bronchial congestion.

PHARMACOLOGY:

Mechanism of action:

Diphenhydramine is an ethanolamines antihistamine, a H1-receptor antagonist. It acts by competing with histamine for H1-receptor sites on the effectors cells, to the exclusion of agonist molecules, without itself initiating a response. This antagonism is competitive and reversible. It does not block the release of histamine. Ammonium chloride is used as an expectorant.

Pharmacokinetic:

Diphenhydramine is readily absorbed from gastrointestinal tract and parental sites of administration. Systemic availability is affected by high first-pass metabolism. Peak plasma concentration is within 1-4 hours after oral administration. It is highly bound to plasma protein, thus widely distributed throughout the body including the CNS. It crosses the placenta and has been detected in breast milk. Most of the diphenhydramine is excreted as metabolites through urine with complete excretion within 24 hours. Ammonium chloride is absorbed from the gastrointestinal tract. The ammonium ion is converted into urea in the liver, the anion thus liberated into the blood stream and extracellular fluid causes a metabolic acidosis and decreases the pH of the urine, this is followed by transient diuresis.

DOSAGE AND ADMINISTRATION:

For oral administration only.

Adult & children > 12 years: 5-10ml, max. 60ml / 24 hours.

Children 6-12 years: 2.5-5ml, max. 30ml / 24 hours.

Children 2-5 years: 2.5ml, max. 15ml / 24 hours.

To be taken 4-6 hourly.

CONTRAINDICATIONS:

Patient who is hypersensitive to diphenhydramine or other antihistamines; patient with MAOIs therapy, impaired renal or hepatic function.

PRECAUTIONS:

Axcel Flemin Expectorant may cause drowsiness and thus do not drive or operate machinery. It should be used with caution in patient with narrow angle glaucoma, history of bronchial asthma, hyperthyroidism, cardiovascular disease, prostatic hypertrophy, peptic ulcer, pyloroduodenal obstructions and bladder neck obstructions. Not to be given to newborns or infants. Not recommended for children below 2 years old. To be used with caution upon the advice of a doctor or pharmacist for children between the age of 2 to 6 years old.

Unsuitable for phenylketonurics.

Pregnancy & lactation:

If the patient is pregnant or nursing, consult doctor before taking this product.

SIDE EFFECTS:

The most common effect is sedation, varying from drowsiness to deep sleep, and including inability to concentrate, lassitude, dizziness, hypotension, muscular weakness, and incoordination. Sedative effects, when they occur, may diminish after a few days. Other side effects include gastrointestinal disturbances, such as nausea, vomiting, diarrhea or constipation, colic and epigastric pain. It may also produce headache, blurred vision, tinnitus, elation or depression, inability, nightmares, anorexia, difficulty in micturition, dryness of the mouth, tightness of chest, tingling, heaviness and weakness of the hands. It may potentiate other CNS depressant compounds. Large doses may precipitate fits in epileptics. The ammonium salts content are irritant to the gastric mucosa and may produce nausea and vomiting particularly in large doses.

DRUG INTERACTIONS:

Additive effects if used with hypnotics, sedatives and tranquilizers. This preparation should not be given to patients receiving MAOIs, aminoglycosides, antibiotics and antimuscarinics. Avoid drinking alcoholic beverages when taking Axcel Flemin Expectorant.

OVERDOSAGE & TREATMENT:

Symptoms of overdosage may vary from CNS depression including sedation, cardiovascular collapse, and stimulation including insomnia, hallucination, tremors or convulsion. Other symptoms include dizziness, tinnitus, ataxia, blurred vision and hypotension. If the drug has been taken recently by mouth, the stomach should be emptied by aspiration and lavage. Emetics are generally of little value. The patient should be kept quiet to minimize the excitation which occurs particularly in children. Convulsion and marked CNS stimulation should preferably be treated with diazepam or phenobarbitone intramuscularly;

paraldehyde, thiopentone sodium, and chlorpromazine have also been recommended. Severe hypotension may require fluid replacement, and assisted respiration may be necessary. Forced diuresis is of little value since antihistamines are rapidly metabolized and only traces are recovered in the urine. Large doses of Ammonium Chloride may cause a profound acidosis and hypokalaemia which should be treated symptomatically.

STORAGE:

Keep container well closed.

Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN

JAUIH DARI KANAK-KANAK

PACK QUANTITIES:

Available in 60ml, 90ml, and 100ml PET bottle; 120ml amber glass bottle.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL19991903AZ

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
No. 1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.