

MICOSTEN-H CREAM

DESCRIPTION:

White smooth homogenous mass

COMPOSITION:

Each gm contains Miconazole Nitrate (Micronised) 20 mg

Hydrocortisone (Micronised) 10 mg

ACTION & PHARMACOLOGY:

Hydrocortisone is a natural corticosteroid with a potent anti-inflammatory and antipruriginous action. Miconazole Nitrate is a synthetic antifungal agent which inhibits the growth of the common dermatophytes and *Candida* sp. together with a strong antibacterial effect against gram-positive bacilli and cocci.

Systemic absorption following topical application to normal skin has not been reported.

INDICATIONS:

Infections of the skin by dermatophytes or *Candida* sp. in which inflammatory symptoms are prominent. Particularly indicated for the initial stages of treatment may be continued with Micosten H Cream. May also be used for mycotic infections with bacterial superinfection.

CONTRAINDICATIONS:

Contraindicated in patients who are sensitive to topical miconazole and hydrocortisone.

Contraindicated in tuberculous skin infections, herpes simplex, vaccinia, all forms of varicella, infections caused by gram-negative bacteria.

SIDE EFFECTS:

Irritation, burning, maceration, allergic contact dermatitis and other sign of skin irritation not present before therapy are some of the isolated reactions reported.

WARNING AND PRECAUTIONS:

Caution is advised with pregnant women and infants under 1 year if the medication is to be applied to extensive surface areas or under occlusive dressing.

Use of topical antifungals may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication. To reduce the possibility of recurrence, treatment should be continued without interruption until the lesion has completely disappeared (usually after 2-5 weeks). Avoid contact with the eyes.

Application to the face should be avoided. Avoid long term continuous topical steroid therapy in infants.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

PREGNANCY AND LACTATION:

It is not known whether topical application is distributed into breast milk. However, problems in humans have not been documented. The cream should not be applied to the breasts prior to nursing.

Elderly:

No geriatric-specific problems have been documented to that to limit the usefulness of this topical application in the elderly. However, it should be used infrequently for brief periods or under close medical supervision in patients with evidence of pre-existing skin atrophy.

DRUG INTERACTIONS:

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and Precautions).

DOSAGE & ADMINISTRATION:

Skin infection: Apply once to twice daily to cover the affected area and rub in gently with finger.

Nail infection: Apply once daily and cover with a bandage.

Treatment should be continued without interruption until the lesion has completely disappeared (usually after 2-5 weeks). After about 7 days of application, if no improvement has occurred, cultural isolation of the offending organism should be followed by appropriate local or systemic anti-microbial therapy.

ROUTE OF ADMINISTRATION:

Topical

OVERDOSE & TREATMENT:

Acute overdosage with the topical application of the drug is unlikely. If irritation or sensitivity reactions occur, use of the medication should be discontinued. In the event of accidental ingestion, an appropriate method of gastric emptying may be used if considered to be necessary.

PRESENTATION:

Aluminium tubes of 5g, 10g, 15g and 50g

STORAGE CONDITION:

Store in a cool dry place below 30°C.

Keep the containers tightly closed.

Protect from light and freezing.

For external use only.

Keep medicine out of reach of children.

MANUFACTURER AND PRODUCT REGISTRATION HOLDER:

Noripharma Sdn.Bhd. 200701034604 (792633-A)

Lot 5030 Jalan Teratai,

5 ½ Miles Off Jalan Meru,

41050 Klang,

Selangor Darul Ehsan, Malaysia.

DATE OF REVISION:

September 2023