



SALBUTAMOL SYRUP

COMPOSITION:

Each 5ml contains
Salbutamol 2mg

Sodium Benzoate 0.2% w/v as preservative.

PRESENTATION:

Colourless syrup with fruit flavor.

INDICATIONS:

For the relief of bronchospasm in asthma, chronic bronchitis and other conditions involving airways obstruction.

PHARMACOLOGY:*Mechanism of Action:*

Salbutamol is a direct-acting sympathomimetic agent with a relatively selective action on beta-2 adrenoceptors. It is used as a bronchodilator in the management of reversible airways obstruction such as asthma and chronic obstructive pulmonary diseases. Salbutamol also decreases uterine contractility and may be given in delaying premature labour.

DOSAGE AND ADMINISTRATION:

Oral administration : 3 or 4 times daily.
Children 2 - 6 years : 2.5 - 5ml (1-2mg).
6 - 12 years: 5ml (2mg).
Over 13 years: 5 - 10ml (2-4mg).

CONTRAINDICATIONS:

Contraindicated in patients with a history of hypersensitivity to any of its component. Salbutamol should not be administered concurrently with beta-blocking drugs, such as propranolol.

PRECAUTIONS:

Caution is required in serious cardiovascular disorder, hyperthyroidism, diabetes mellitus, closed-angle glaucoma and hypertension. Prolonged use in high dosage should be avoided as resistance may develop. Hypokalaemia is known to be a possible side-effect during treatment with salbutamol and this may be enhanced during concomitant therapy of corticosteroid, diuretics or theophylline and serum potassium should be monitored in these circumstances.

Use in Pregnancy and Lactation:

Unnecessary administration of Salbutamol during the first trimester of pregnancy is undesirable.

Newborn / Premature:

No special precaution stated for administration to newborn and premature infants.

SIDE EFFECTS:

The most common side effect is fine tremor of skeletal muscle (particularly the hands), palpitations, and muscle cramps. Hyperactivity in children. Slight tachycardia, tenseness, headaches, and peripheral vasodilation have been reported after large doses. Potentially serious hypokalaemia may result from beta-2 agonist therapy.

DRUG INTERACTION:

An enhanced hypokalaemia effects may occur during co-administration with diuretics, theophylline, steroids and muscle relaxants.

OVERDOSAGE AND TREATMENT:

Symptoms of over dosage generally include; tachycardia, CNS stimulation, tremor, hypokalaemia, and hyperglycaemia. Symptomatic treatment of the adverse effects has proved successful.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in 60ml & 100ml PET bottle.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL 19992588AZ



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (00082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.
Tel +606 336 2222

Regn - 060514