

FIBID CAPSULE

DESCRIPTION:

Size '0' scarlet/white coloured hard gelatin capsule containing white to off white powder.

COMPOSITION:

Each capsule contains Gemfibrozil 300mg

ACTIONS & PHARMACOLOGY:

Gemfibrozil is a lipid regulating agent which reduces total cholesterol, LDL cholesterol, VLDL and triglycerides and increases HDL cholesterol. Gemfibrozil's mechanism of action has not been definitely established. In man, it inhibits peripheral lipolysis and decreases the hepatic extraction of free fatty acids. It also inhibits synthesis and increases clearance of apolipoprotein B which is a carrier of VLDL. It increases the HDL₂ and HDL₃ subfractions as well as apolipoprotein A-I and A-II. Animal studies suggest that the turnover and removal of cholesterol from the liver is increased.

Gemfibrozil is well absorbed from the GIT after oral administration. Peak plasma levels occur in 1-2 hours with plasma half-life of 1.5 hours following single dose and 1.3 hours following multiple doses. Plasma levels appear proportional to dose and do not demonstrate accumulation across time following multiple doses. It undergoes oxidation of a ring methyl group to form successively a hydroxymethyl and a carboxyl metabolite. Approximately 70% of administered dose is excreted in the urine as the glucuronide conjugate and as unchanged drug. About 6% of the administered dose is excreted in the faeces.

INDICATIONS:

Primary prevention of coronary heart disease in patients with hypercholesterolemia, mixed dyslipidemia and hypertriglyceridemia. Fredrickson's classification Types IIa, IIb, IV respectively who have not responded adequately to diet and other appropriate measures. Treatment of adult patients with elevated levels of serum triglycerides (types IV and V hyperlipidemia) who present a risk of pancreatitis and who do not respond adequately to a determined dietary effort to control them.

CONTRAINDICATIONS:

Contraindicated in patients who have previously exhibited hypersensitivity to it. Also contraindicated in patients with primary biliary cirrhosis, gallbladder disease or gallstones, severe renal and hepatic function impairment.

SIDE EFFECTS:

The most common adverse effects are gastrointestinal complaints (dyspepsia, nausea, vomiting, anorexia, diarrhoea). Headache, dizziness, visual disturbances, allergic skin reactions, muscular and joint pains, anaemia and leukopenia have also been reported. In a few cases an increase in biliary lithogen index was observed. Rarely, elevated SGOT, SGPT, serum alkaline phosphatase, LDH and bilirubin values may occur, but these are usually reversible after discontinuation of gemfibrozil therapy.

PRECAUTIONS:

If patients receiving Gemfibrozil experience muscle pain or weakness, evaluation for myositis is recommended. It is recommended that Gemfibrozil be withdrawn if myositis is diagnosed. Gemfibrozil therapy may be discontinued,

- (a) If gallstones are found.
- (b) If results of hepatic function tests show significant abnormalities.
- (c) If response is inadequate after three months of treatment.

PREGNANCY & LACTATION:

Gemfibrozil has not been studied in pregnant women; hence safety in pregnancy and nursing mothers has not been established.

DRUG INTERACTION:

Concurrent use of anticoagulants may increase the effect of the anticoagulant. The anticoagulant dose should be reduced to maintain the prothrombin time at the desired level to prevent bleeding complications. Concurrent use of Lovastatin (or other HMG-CoA reductase inhibitors) may cause severe myositis and myoglobinuria.

DOSAGE & ADMINISTRATION:

The recommended daily dose is 900-1200mg. The maximum daily dose is 1500mg. Adult: Oral, 1.2gm a day in two divided doses thirty minutes before the morning and evening meals. Pediatric: Dosage has not been established.

ROUTE OF ADMINISTRATION:

Oral administration

OVERDOSAGE & TREATMENT:

While there has been no reported case of overdosage, symptomatic and supportive measures should be taken if it occurs.

PRESENTATION:

Blister of 10 capsules. Box of 30, 100, 250, 500 and 1000 capsules.

STORAGE:

Store in a cool dry place below 30°C. Keep medicine out of reach of children. Protect from light.

SHELF-LIFE:

3 years

MANUFACTURER AND PRODUCT REGISTRATION HOLDER:

NORIPHARMA SDN. BHD. (792633-A)
Lot 5030, Jalan Teratai, 5 1/2 Mile off Jalan Meru,
41050 Klang, Selangor Darul Ehsan, Malaysia.

Revision Date: July 2020